

*International Europhysics Conference on High
Energy Physics, 2007*

Charmless Hadronic B Decays at *BABAR*

— July 20 —

J. P. Burke



UNIVERSITY OF
LIVERPOOL

for the *BABAR* collaboration

Outline of talk

- Quick intro to charmless hadronic B decays, overview of $BABAR$ detector, common analysis techniques
- Preliminary $BABAR$ results:
 - New *previously unmeasured* ($K^+ K^- \pi^+$)/*unstudied (the rest)*:

$B^+ \rightarrow K^+ K^- \pi^+$ *Even number of kaons in final state*

$B^+ \rightarrow a_1(1260)\pi$ *$B \rightarrow AP$, rather than well covered $B \rightarrow PP, VP$*

$B \rightarrow b_1(1235)h^+$ *Also $B \rightarrow AP$, G -parity suppression*


$B^+ \rightarrow \eta_X K^+$ *Possible η, η' excitations, no definitive theory*

- Updated:

$B \rightarrow K\pi^0, \pi\pi^0$ *$B \rightarrow K\pi$ excellent probe for new physics*

$B \rightarrow h_1 h_2$ ($h_1 = \eta, \eta', \omega$; $h_2 = K^+, \pi^+, K^0$), followed by summary


General overview of charmless hadronic B decays





- Interfering SM amplitudes ... good place to look for **direct CPV**—see **Nicolas Arnaud's** talk
- Ideal environments to study loop processes (where **new physics** may enter)
- Extract **CKM parameters**—see talks by **Mark Allen**, **Josh Thompson**, **Emmanuel Latour** (c.f. $b \rightarrow c$ results to constrain NP)
- Use measured rates phenomenologically to test/develop theoretical models (**factorization**, **pQCD**, **SU(3) flavour symmetry**, ...)

The experiment

PEP-II collider: $9 \text{ GeV } e^-$, $3.1 \text{ GeV } e^+$,
 data taking commenced 1999



$\approx 232 \times 10^6 B\bar{B}$ pairs




$\approx 383 \times 10^6 B\bar{B}$ pairs

(the rest)

BABAR Detector

- Muon/Hadron Detector
- Magnet Coil
- Electron/Photon Detector
- Cherenkov Detector
- Tracking Chamber
- Support Tube
- Vertex Detector



Common analysis strategy

- Aim is to isolate **very small signal** from **vast backgrounds**
- Main background due to **continuum** events, $e^+e^- \rightarrow d\bar{d}, u\bar{u}, s\bar{s}, c\bar{c}$
- ($\#$ signal events):($\#$ continuum events) enhanced using
 - **Particle ID** systems
 - **Cutting** on discriminating event variables
- Discriminating variables include
 - Kinematic: $m_{ES}, \Delta E$ —use beam and decay products' (E, \vec{p})
 - Event shape: B events isotropic, continuum events jet-like
 - Combine in **Fisher discriminants**, \mathcal{F} , and **neural networks**, \mathcal{N}
 - Resonance: **invariant mass**, **helicity angle** (related to **spin**)
- Signal parameters extracted using **maximum likelihood (ML)**
- Must also treat background from B events

$B^+ \rightarrow K^+ K^- \pi^+$ —preliminary

- Heavily suppressed $B^+ \rightarrow \bar{K}^{*0} K^+$ expected to occur via $b \rightarrow d$ penguin, no tree
- Search conducted with no significant signal observed:

$$\mathcal{B}(B^+ \rightarrow \bar{K}^{*0}(892)K^+) < 1.1 \times 10^{-6} \dagger$$

90% CL

$$\mathcal{B}(B^+ \rightarrow \bar{K}_0^{*0}(1430)K^+) < 2.2 \times 10^{-6}$$

arXiv:0706.1059 [hep-ex]

... in agreement with pQCDF and SU(3) predictions

\dagger : assists in bounding $|\sin 2\beta(B^0 \rightarrow J/\psi K_S^0) - \sin 2\beta_{eff}(B^0 \rightarrow \phi K_S^0)| < 0.11$ (in SM)


- $\mathcal{B}(B^+ \rightarrow \phi \pi^+) < 2.4 \times 10^{-7}$ (also $b \rightarrow d$ penguin, need *three* gluons or Z^0/γ from loop) arXiv:hep-ex/0605037
- Other (known, charmless) possible contributions to Dalitz plot include f_0 's and non-resonant ... not expected to be very large
- Enough for a significant signal in the inclusive decay? ...

$B^+ \rightarrow K^+ K^- \pi^+$ — preliminary


- ML technique applied to full Dalitz plot using m_{ES} , ΔE , \mathcal{N}
- Good sized (429 events) signal observed:

$$\mathcal{B}(B^+ \rightarrow K^+ K^- \pi^+) = (5.0 \pm 0.5 \pm 0.5) \times 10^{-6} \quad 9.6\sigma$$


- Dominated by broad structure in $K^+ K^-$ spectrum $\sim 1.5 \text{ GeV}/c^2$
- Similar structure seen in DP analyses of $B^+ \rightarrow K^+ K^- K^+$, $B^0 \rightarrow K^+ K^- K^0$ (not in $K^\pm K^0$ spectrum, not in D decays)—dubbed $X_0(1550)$ by *BABAR*, $f_X(1500)$ by Belle



ΔE proj. plot



DP s Plot




$m_{K^+K^-}$ s Plot

$B^+ \rightarrow a_1(1260)\pi$ —*preliminary*

- First evidence of $B^+ \rightarrow a_1^+(1260)\pi^0$ and $B^+ \rightarrow a_1^0(1260)\pi^+$ ($b \rightarrow u\bar{u}d$ tree + penguin), final state $\pi^+\pi^-\pi^+\pi^0$
- ML using m_{ES} , ΔE , \mathcal{F} , invariant mass of reconstructed a_1 , helicity angle; results consistent with factorization predictions
- With $\mathcal{B}(a_1^+(1260) \rightarrow 3\pi^\pm) = \frac{1}{2}$, $\mathcal{B}(a_1^0(1260) \rightarrow \pi^+\pi^-\pi^0) = 1$:

$$\mathcal{B}(B^+ \rightarrow a_1^+(1260)\pi^0) = (26.4 \pm 5.4 \pm 4.1) \times 10^{-6} \quad 4.2\sigma$$

$$\mathcal{B}(B^+ \rightarrow a_1^0(1260)\pi^+) = (20.4 \pm 4.7 \pm 3.4) \times 10^{-6} \quad 3.8\sigma$$



← Projection plots of m_{ES} (likelihood cut applied)

- Helpful for measurement of UT angle α from $\rho\pi$

$B \rightarrow b_1(1235)h^+ \text{---preliminary}$

- Recent searches of $B \rightarrow A\pi$ have revealed rather large \mathcal{B} 's, e.g. $\mathcal{B}(B^0 \rightarrow a_1^\pm(1260)\pi^\mp) = (33.2 \pm 3.8 \pm 3.0) \times 10^{-6}$ PRL 97, 051802, and $B^+ \rightarrow a_1(1260)\pi$ as shown on previous slide
- Two types of axial-vector mesons, A :
 - a_1 is the $I^G = 1^-$ member of the $J^{PC} = 1^{++}$ 3P_1 nonet ($\uparrow\uparrow$)
 - b_1 is the $I^G = 1^+$ member of the $J^{PC} = 1^{+-}$ 1P_1 nonet ($\uparrow\downarrow$)
- K_{1A} (K_{1B}) member of 3P_1 (1P_1) nonet, K_{1A} and K_{1B} mix to give physical $K_1(1270)$ and $K_1(1400)$
- Mixing angle θ known (within a few $^\circ$'s) up to twofold ambiguity symmetric about $\frac{\pi}{4}$ (32° and 58°)
- From naïve factorization (for non-G-parity-suppressed modes)
 - hep-ph/0602243 v4: $\mathcal{B}(B \rightarrow b_1 h^+) \approx 4 - 30 \times 10^{-6}$ with $\theta = 32^\circ$, $\mathcal{B}(B \rightarrow b_1 h^+) \approx 4 - 20 \times 10^{-7}$ with $\theta = 58^\circ$
 - arXiv:0705.1181 [hep-ph]: $\mathcal{B}_{32^\circ}(B \rightarrow b_1 h^+) \approx 18 - 36 \times 10^{-6}$

$B \rightarrow b_1(1235)h^+ \text{—preliminary}$

- $b \rightarrow u\bar{u}q_d$ ($q_d = d, s$) tree (K Cabibbo suppressed wrt π) + penguin (π suppressed wrt K)
- b_1 reco'd as $b_1 \rightarrow \omega(\rightarrow \pi^+\pi^-\pi^0(\rightarrow \gamma\gamma))\pi$
- ML: $m_{ES}, \Delta E, \mathcal{F}$, invariant masses of reconstructed b_1 and ω
- **First observations** of $B \rightarrow b_1(1235)h^+$ decays:

$\mathcal{B}(B^+ \rightarrow b_1^0(1235)\pi^+) = (6.6 \pm 1.7 \pm 1.0) \times 10^{-6}$	4.0 σ	<u>$b\pi \Rightarrow \theta$</u>
$\mathcal{B}(B^+ \rightarrow b_1^0(1235)K^+) = (8.8 \pm 1.7 \pm 1.0) \times 10^{-6}$	5.3 σ	<u>$= 32^\circ$</u> ;
$\mathcal{B}(B^0 \rightarrow b_1^\mp(1235)\pi^\pm) = (10.9 \pm 1.2 \pm 0.9) \times 10^{-6} \dagger$	8.9 σ	... <u>bK in</u>
$\mathcal{B}(B^0 \rightarrow b_1^-(1235)K^+) = (7.4 \pm 1.0 \pm 1.0) \times 10^{-6}$	6.1 σ	<u>between</u>


\dagger : Asymmetry measurement: $b_1^+\pi^-$ heavily suppressed ($(0 \pm 10)\%$) wrt $b_1^-\pi^+$ — follows (in SM) from b_1 decay const. ~ 0 (a prediction of G-parity suppression)

Best agreement: hep-ph/0602243 (32°), but b_1K \mathcal{B} 's smaller than predicted

$B \rightarrow b_1(1235)h^+ \text{---preliminary}$

Top row: $B^+ \rightarrow b_1^0(1235)\pi^+$; second row: $B^+ \rightarrow b_1^0(1235)K^+$;

Prelim.




third row: $B^0 \rightarrow b_1^\mp(1235)\pi^\pm$; bottom row: $B^0 \rightarrow b_1^-(1235)K^+$

$$B^+ \rightarrow (\eta_X)_{1.2-1.8 \text{ GeV}/c^2} K^+ \text{—preliminary}$$

- B decays to a kaon and excited states of η , η' ($b \rightarrow s$ penguin)
- Cands. for such excited $J^P = 0^-$ states $\eta(1295)$, $\eta(1405)$, $\eta(1475)$
- Dynamics of these quasi-2-body decays is a difficult theoretical problem ... why is $\mathcal{B}(B^+ \rightarrow \eta' K^+)$ so large?
- Explanation thought to be in SM, but large uncertainties (higher-order corrections/charming-penguin contributions to factorization, SCET, exotic gluonium states ...?)
- η_X 's reco'd through decays to $K_S^0 K^\pm \pi^\mp$ (via $K^* \bar{K}$) and $\eta \pi^+ \pi^-$ in invariant mass ranges $1.2 - 1.8$, $1.2 - 1.5 \text{ GeV}/c^2$
- Spectra here not well known *J. Phys. G33, 1 (July 2006)—p591*
- $f_1(1285)$ and $f_1(1420)$ ($J^P = 1^+$), and $\phi(1680)$ ($J^P = 1^-$), also found in these spectra—these states are also considered

$$B^+ \rightarrow (\eta_X)_{1.2-1.8 \text{ GeV}/c^2} K^+ \text{—preliminary}$$

- Two ML fits to m_{ES} , ΔE , \mathcal{F} , invariant η_X mass, helicity angle
- First fit simultaneously extracts \mathcal{B} 's for $B^+ \rightarrow \eta_X (\rightarrow K_S^0 K^\pm \pi^\mp) K^+$, where $\eta_X = \eta(1405), f_1(1420), \eta(1475), \phi(1680)$
- Second fit simultaneously extracts \mathcal{B} 's for $B^+ \rightarrow \eta_X (\rightarrow \eta \pi^+ \pi^-) K^+$, where $\eta_X = f_1(1285), \eta(1295), \eta(1405), f_1(1420)$
- m_{ES} , ΔE and \mathcal{F} discriminate signal versus background
- Invariant η_X mass and helicity angle discriminate between signal hypotheses (different masses, widths, spins)



$$B^+ \rightarrow (\eta_X)_{1.2-1.8 \text{ GeV}/c^2} K^+ \text{—preliminary}$$

$$\mathcal{B}(\eta_X \rightarrow K^* \bar{K}) \times$$

$\mathcal{B}(B^+ \rightarrow \eta(1475)K^+) = (13.8 \pm 1.8 \pm 1.0) \times 10^{-6} \dagger$	$7.5\sigma \dagger$
$\mathcal{B}(B^+ \rightarrow \eta(1405)K^+) < 1.2 \times 10^{-6}$	90% CL
$\mathcal{B}(B^+ \rightarrow f_1(1420)K^+) < 4.1 \times 10^{-6}$	90% CL
$\mathcal{B}(B^+ \rightarrow \phi(1680)K^+) < 3.4 \times 10^{-6}$	90% CL

\dagger : only if only S-wave contributions are signal, phase-space $K^* K K$

$$\mathcal{B}(\eta_X \rightarrow \eta \pi \pi) \times$$

$\mathcal{B}(B^+ \rightarrow \eta(1295)K^+) = (2.9 \pm 0.8 \pm 0.2) \times 10^{-6}$	3.5σ
$\mathcal{B}(B^+ \rightarrow \eta(1405)K^+) < 1.3 \times 10^{-6}$	90% CL
$\mathcal{B}(B^+ \rightarrow f_1(1285)K^+) < 0.8 \times 10^{-6}$	90% CL
$\mathcal{B}(B^+ \rightarrow f_1(1420)K^+) < 2.9 \times 10^{-6}$	90% CL

- Same mechanism at work as for $B^+ \rightarrow \eta' K^+$?

Updated measurements—preliminary

- $B^0 \rightarrow K^0 \pi^0$ \mathcal{B} updated: $(10.34 \pm 0.66 \pm 0.58) \times 10^{-6}$
 - $K^0 \rightarrow K_S^0$ used for updated time-dependent CP asymmetry measurements, including $\sin 2\beta_{eff}$ —see Josh Thompson's talk
- $B \rightarrow h\pi^0$ \mathcal{B} 's updated—these modes can be used to constrain α , see Mark Allen's talk
 - $\mathcal{B}(B^0 \rightarrow \pi^0 \pi^0) = (1.47 \pm 0.25 \pm 0.12) \times 10^{-6}$
 - $\mathcal{B}(B^+ \rightarrow \pi^+ \pi^0) = (5.02 \pm 0.46 \pm 0.29) \times 10^{-6}$
 - $\mathcal{B}(B^+ \rightarrow K^+ \pi^0) = (13.6 \pm 0.6 \pm 0.7) \times 10^{-6}$

Lipkin ratio consistent with 1:

$$\begin{aligned}
 R_L &= \frac{2\Gamma(B^+ \rightarrow K^+ \pi^0) + 2\Gamma(B^0 \rightarrow K^0 \pi^0)}{\Gamma(B^+ \rightarrow K^0 \pi^+) + \Gamma(B^0 \rightarrow K^+ \pi^-)} = 1 + \mathcal{O}\left(\frac{P_{EW} + T}{P}\right)^2 \\
 &= \boxed{1.11 \pm 0.07}
 \end{aligned}$$

Updated measurements

- $B \rightarrow h_1 h_2$ ($h_1 = \eta, \eta', \omega$; $h_2 = K^+, \pi^+, K^0$):

$$\mathcal{B}(B^+ \rightarrow \eta\pi^+) = (5.0 \pm 0.5 \pm 0.3) \times 10^{-6}$$

$$\mathcal{B}(B^+ \rightarrow \eta K^+) = (3.7 \pm 0.4 \pm 0.1) \times 10^{-6}$$

$$\mathcal{B}(B^+ \rightarrow \eta'\pi^+) = (3.9 \pm 0.7 \pm 0.3) \times 10^{-6}$$

$$\mathcal{B}(B^+ \rightarrow \eta'K^+) = (70.0 \pm 1.5 \pm 2.8) \times 10^{-6}$$

$$\mathcal{B}(B^0 \rightarrow \eta'K^0) = (66.6 \pm 2.6 \pm 2.8) \times 10^{-6}$$

$$\mathcal{B}(B^+ \rightarrow \omega\pi^+) = (6.7 \pm 0.5 \pm 0.4) \times 10^{-6}$$

$$\mathcal{B}(B^+ \rightarrow \omega K^+) = (6.3 \pm 0.5 \pm 0.3) \times 10^{-6}$$

$$\mathcal{B}(B^0 \rightarrow \omega K^0) = (5.6 \pm 0.8 \pm 0.3) \times 10^{-6}$$

- See [Nicolas Arnaud's](#) talk for asymmetry measurements

-
- Further charmless results presented by [Silvano Tosi](#): $B \rightarrow p\bar{p}h$,

$$B \rightarrow \phi K^*$$

Summary

- $B^+ \rightarrow K^+ K^- \pi^+$: first observation of charmless 3-body B meson decay to final state with even number of kaons
- $B \rightarrow Ah$ ($A =$ axial-vector mesons b_1, a_1 ; $h = K, \pi$): first observation of three modes, evidence of a further three modes
- First observation of G-parity suppression in B decays ($B^0 \rightarrow b_1^+(1235) \pi^-$)
- Excess of events that could be $B^+ \rightarrow \eta(1475)K^+$, $B^+ \rightarrow \eta(1295)K^+$... are these η, η' excitations?
- Value of Lipkin ratio updated
- Several previous results updated with greater precision
- New/improved limits placed on numerous modes

Backup slides . . .


Reminder of some winter conference results

- $\mathcal{B}(B^+ \rightarrow \rho^+ K^0) = \underline{(8.0 \pm 1.4 \pm 0.6) \times 10^{-6}} \quad 7.9\sigma$
 - First observation of this pure penguin mode [hep-ex/0702043](#)—
useful for determining UT angle γ using U-spin and charmless $B^+ \rightarrow M^+ M^0$ decays [Phys. Lett. B635, 330](#)
- $\mathcal{B}(B^0 \rightarrow K_1^+(1270)\pi^-) = \underline{(12.0 \pm 3.1 \pm 9.3)[< 25.2] \times 10^{-6}} \quad 2.3\sigma$
 $\mathcal{B}(B^0 \rightarrow K_1^+(1400)\pi^-) = \underline{(16.7 \pm 2.6 \pm 5.0)[< 21.8] \times 10^{-6}} \quad 3.0\sigma$
 - Needed to pin down penguin amplitudes in order to extract α from $B^0 \rightarrow a_1^\pm(1260)\pi^\mp$


Topological variables

- $B\bar{B}$ pairs produced almost at rest in CM frame; $q\bar{q}$ pairs from continuum have considerable KE

Isotropic B event



Cut here, for example




- Can combine variables in Fisher discriminants, \mathcal{F} , and neural networks, \mathcal{N}


Kinematic variables

- Want to optimise resolution, take full advantage of available info, minimise correlations, allow for asymmetric nature of collider:

$$m_{ES} = \sqrt{E_{beam}^{*2} - \vec{p}_B^{*2}}$$







$$\Delta E = E_B^* - E_{beam}^*$$



- E_{beam}^* is expected energy of reco'd candidate using beam's *4-momentum* and detected decay products' *3-momenta* (in lab)
- ΔE has inferior resolution but since uses detected decay products' mass hypotheses is sensitive to particle mis-ID

Resonance variables

- For decays via intermediate state(s) can utilise resonance **invariant mass** and **helicity angle**, θ_H



- Definition of θ_H depends on $\#$ of resonance daughters
- E.g., for two daughters, it's the polar angle in the resonance's rest frame of one of its daughters where the polar axis is anti-parallel to the receding B frame

The maximum likelihood (ML) method

- **Probability density functions (PDFs)**, with parameters $\vec{\theta}$, are used to describe distributions of event variables \vec{x} (e.g. m_{ES} , ΔE , \mathcal{F} , ...) for various species (e.g. signal, continuum background)
- ML technique used to determine values of $\vec{\theta}$ that **maximise** the probability of obtaining the observed measurements according to the pre-determined PDF forms (e.g. Gaussian)
- For the *extended* ML, the values of \vec{n} are also extracted where \vec{n} are the yields for each of our species
- The probability to maximise is given by the **extended likelihood function** ...

The extended likelihood function

- The normal likelihood is the product of the PDFs for each individual candidate i (of which there are N), $\prod P(\vec{x}_i; \vec{\theta})$
- The extended likelihood function is given by $\mathcal{L}(\vec{n}, \vec{\theta}) = \text{Poisson factor} \times \text{normal likelihood function}$
- Omitting constants the function to maximise is

$$\mathcal{L}'(\vec{n}, \vec{\theta}) = \exp\left(-\sum_{k=1}^M n_k\right) \prod_{i=1}^N \left(\sum_{j=1}^M n_j \left(\prod_{l=1}^V \mathcal{P}_j^l(x_i^l; \vec{\theta})\right)\right)$$

- The PDF $P(\vec{x}_i; \vec{\theta})$, for a given measurement i , is the sum of the PDFs for each hypothesis (of which there are M)
- The PDF for each hypothesis, \mathcal{P}_j , is the product of the individual PDFs, \mathcal{P}^l , for each of the V discriminating event variables (assuming negligible correlation, which must be shown)

sPlots

- Used to reconstruct a variable distribution for a particular species (e.g. signal) from the PDFs of other (fit) variables, \mathcal{P}
- sWeight for species of interest assigned to each event:

$$\frac{\sum_{j=1}^{N_S} \mathbf{V}_{nj} \mathcal{P}_j}{\sum_{k=1}^{N_S} N_k \mathcal{P}_k}$$

- N_S is # of species, \mathbf{V} is the covariant matrix from the fit, N_k is the yield for species k returned from the fit, subscript n refers to the species of interest
- Summing the sWeights over all events gives the species yield
- Bin each sWeighted event to reproduce (e.g. signal) distribution —i.e. signal *sPlot*—of that variable (weight only the variable that we're *sPlotting*)

Dalitz plot analyses

- Note, *amplitude-level Dalitz plot* analyses are beginning to dominate charmless 3-body decays as datasets \uparrow (but none here!) (see [Josh Thompson's](#) presentation for a discussion of the $B^0 \rightarrow K^+ K^- K^0$ DP analysis)
- Structure in the DP gives info on resonance masses, widths, spins, *relative phases*, *interference*
- Model each contribution to the DP as a separate amplitude with a complex coefficient (*isobar* model)

