



**ICHEP 2002**

# Direct CP asymmetries and charmless branching fractions with BABAR

**Adrian Bevan**




THE UNIVERSITY  
*of* LIVERPOOL

**For the BABAR Collaboration**



# Overview

- Motivation: Direct CP Violation (CPV)  
New Physics sensitivity e.g.  $X_S\gamma$ ,  $\phi_K$   
Penguins



- Event Selection Methods
- Results: Charmless decays,  $K^*\gamma$ ,  $D^0_{CP}K^-$
- Summary and outlook



# Direct CP Violation

$$|A_f|^2 - |\bar{A}_{\bar{f}}|^2 \neq 0 \quad \longrightarrow \quad \text{direct CPV}$$

- need interference between diagrams with different strong ( $\delta_i$ ) and weak phases ( $\phi_i$ ):

$$\longrightarrow |A_f|^2 - |\bar{A}_{\bar{f}}|^2 = -2 \sum_{i,j} A_i A_j \sin(\phi_i - \phi_j) \sin(\delta_i - \delta_j)$$

- Direct CPV **only** seen in  $K^0 \bar{K}^0$ ;  $\epsilon' \sim \text{few } 10^{-6}$
- theory predicts large asymmetries in  $B^{+0}$  (**few to 80%**)
- Diagrams from New Physics (e.g. SUSY) can modify SM asymmetries/branching fractions



• Experimentally - look for CP asymmetries:

$$A_{CP} = \frac{Br(\bar{B} \rightarrow \bar{f}) - Br(B \rightarrow f)}{Br(B \rightarrow f) + Br(\bar{B} \rightarrow \bar{f})}$$

• Neutral B – time dependent asymmetry

$$f_{\pm}(\Delta t) = \frac{e^{-|\Delta t|/\tau}}{4\tau} \left[ 1 \pm S_f \sin(\Delta m_d \Delta t) \mp C_f \cos(\Delta m_d \Delta t) \right]$$

direct CPV

**C ≠ 0** ← direct CPV; e.g. for  $\rho\pi$ ,  $\pi\pi$  ...

• Charged B – time integrated asymmetry (no mixing); also used for some  $B^0$  results (e.g.  $K^0\pi^0$ )




# Searching for direct CPV

- Large  $A_{CP}$  requires amplitudes of similar order
  - $b \rightarrow u$ : suppressed tree: charmless decays
    - large predicted  $A_{CP}$
  - $b \rightarrow s, d$ : penguins: radiative decays
    - small predicted  $A_{CP}$

- Understand penguins

- Access to  $\alpha$  and  $\gamma$



- Sensitive to New Physics effects via loops

- minimal SUGRA:  $B \rightarrow X_s \gamma, \phi K^+, K^0 \pi^+ \dots$
- R-parity Violating SUSY:  $\phi K_s \dots$
- SUSY searches –  $K^* \gamma$



# Experimental Issues

- small branching fractions
  - large continuum background (u,d,s,c)
  - other B background
- charge bias
  - detector: trigger, tracking; reconstruction
  - event selection, particle ID, analysis

small  
systematic  
error:

- Proof of principle

- $A_{CP}$  in  $J/\psi\pi$ ,  $J/\psi K$  PRD 65 091101 (2002)

- $A_{J/\psi\pi} = 0.01 \pm 0.22 \pm 0.01$

- $A_{J/\psi K} = 0.003 \pm 0.030 \pm 0.004$

few per mille





# Event Selection Techniques

Use beam energy to constrain mass & energy

$$\Delta E = E_B - E_{beam}^*$$

$$m_{ES} = \sqrt{(E_{beam}^*)^2 - P_B^2}$$

$\Delta E$   $\sigma \sim 15\text{-}80$  MeV; larger with neutrals







# Event Selection Techniques

→ B events are spherical

→ u,d,s,c is jet-like



- Fisher Discriminants
- Thrust
- Sphericity
- Fox-Wolfram moments
- flavour-tagging (e,  $\mu$ , K, slow  $\pi$  from other B)



- Maximum Likelihood fits or cut based analysis
- off-resonance &  $\Delta E$  sidebands are used to parameterise light quark background



# RESULTS





# Two body decays

can measure

$\alpha$  eg.  $B \rightarrow \pi^+ \pi^-$ ,  $\pi^+ \pi^0$ ,  $\pi^0 \pi^0$  see Jim Olsen's talk

$\gamma$  eg.  $B \rightarrow K^+ \pi^-$ ,  $K^0 \pi^+$  using Fleischer Mannel bound

$A_{CP}$  can be sizeable



Plots have an optimised cut on likelihood ratio



Preliminary

New results →

~88 × 10<sup>6</sup> B pairs

~60 × 10<sup>6</sup> B pairs

Mode	N <sub>EVENTS</sub>	Branching ratio (×10 <sup>-6</sup> )	A <sub>CP</sub>
B <sup>0</sup> → K <sup>+</sup> π <sup>-</sup>	589 ± 30	(17.9 ± 0.9 ± 0.6)	-0.102 ± 0.050 ± 0.016
B <sup>+</sup> → K <sup>+</sup> π <sup>0</sup>	239 ± 22 ± 6	(12.8 ± 1.2 ± 1.0)	-0.09 ± 0.09 ± 0.01
B <sup>+</sup> → π <sup>+</sup> π <sup>0</sup>	125 <sup>+23</sup> <sub>-21</sub> ± 10	(5.5 ± 1.0 ± 0.6)	-0.03 ± 0.18 ± 0.02
B <sup>0</sup> → K <sup>0</sup> π <sup>0</sup>	86 ± 13 ± 3	(10.4 ± 1.5 ± 0.8)	0.03 ± 0.36 ± 0.09
B <sup>0</sup> → K <sup>+</sup> K <sup>-</sup>	< 15.9	< 0.6 (90% C.L.)	-
B <sup>+</sup> → K <sup>+</sup> K <sup>0</sup>	< 10	< 1.3 (90% C.L.)	-
B <sup>+</sup> → K <sup>0</sup> π <sup>+</sup>	172 ± 17 ± 9	(17.5 ± 1.8 ± 1.3)	-0.17 ± 0.10 ± 0.02

← 5% ΔA<sub>CP</sub>



hep-ex/0206053

All upper limits are @ 90% C.L.



# $B \rightarrow hhh, h = \pi, K$

## Motivation

- measure  $\gamma$  using a full Dalitz plot
- look for direct CPV in charge asymmetry

## Method



- cut based analysis across the Dalitz plot for branching ratio:

$$Br(B \rightarrow f) = \frac{1}{N_{B\bar{B}}} \sum_i \frac{(N_{OBSERVED} - N_{udsc} - N_{bkgd})}{\epsilon_i}$$



Diagram annotations:

- data**: points to the  $N_{OBSERVED}$  term.
- light q background**: points to the  $N_{udsc}$  term.
- B background**: points to the  $N_{bkgd}$  term.
- bin efficiency**: points to the  $\epsilon_i$  term.
- sum over Dalitz plot**: points to the summation index  $i$ .

- B backgrounds:  $J/\psi K, D\pi/DK, X\text{-feed}; D \rightarrow \pi\pi, K\pi, KK$ 
  - subtract open charm from  $J/\psi, \psi_{(2s)}$  and  $D^0$
- measure  $Br(B^- \rightarrow D^0 \pi^-)$  as a cross check



Preliminary





## Branching fractions measured across the Dalitz plot

$$Br(B^\pm \rightarrow K^\pm \pi^\mp \pi^\pm) = (59.2 \pm 4.7 \pm 4.9) \times 10^{-6}$$

$$Br(B^\pm \rightarrow K^\pm K^\mp K^\pm) = (34.7 \pm 2.0 \pm 1.8) \times 10^{-6}$$

$$Br(B^\pm \rightarrow \pi^\pm \pi^\mp \pi^\pm) < 15 \times 10^{-6} \quad (90\% \text{ C.L.})$$

$$Br(B^\pm \rightarrow K^\pm K^\mp \pi^\pm) < 7 \times 10^{-6} \quad (90\% \text{ C.L.})$$

### • Main systematic uncertainties

- PID 3 ~ 6%
- tracking 2.4%



# $B \rightarrow \eta' K^{(*)}, \omega K, \omega \pi^{\pm}$

- $B \rightarrow \eta' K^{(*)}$ , suppressed tree;  $A_{CP}$  4~30% (theory) **sensitive to  $\gamma$**
- $B \rightarrow \omega K, \omega \pi^{\pm}$ , penguin;  $A_{CP} \sim 6\%$  (theory) **sensitive to  $\alpha, \gamma$**

$\sim 60 \times 10^6 B$  pairs preliminary

$Br(B^{\pm} \rightarrow \eta' K^{\pm}) = (67 \pm 5 \pm 5) \times 10^{-6}$

$Br(B^0 \rightarrow \eta' K^0) = (46 \pm 6 \pm 4) \times 10^{-6}$

$Br(B^0 \rightarrow \eta' K^{*0}) < 13 \times 10^{-6}$


$\sim 22.7 \times 10^6 B$  pairs

$Br(B^0 \rightarrow \omega \pi^{\pm}) = (6.6^{+2.1}_{-1.8} \pm 0.7) \times 10^{-6}$

$\sim 22.7 \times 10^6 B$  pairs


$A_{CP}(B^{\pm} \rightarrow \eta' K^{\pm}) = -0.11 \pm 0.11 \pm 0.02$

$A_{CP}(B^{\pm} \rightarrow \omega \pi^{\pm}) = -0.01^{+0.29}_{-0.31} \pm 0.03$





Preliminary



$\sim 60 \times 10^6$  B pairs


First Observation!

Signal yield =  $26.6^{+7.7}_{-6.6}$  events

$$Br(B^0 \rightarrow \omega K^0) = (5.9^{+1.7}_{-1.5} \pm 0.9) \times 10^{-6}$$

6.6  $\sigma$  significance  
(statistical)

- penguin
- sensitive to  $\alpha, \gamma$
- expect small  $A_{CP}$
- ML fit
- dominant systematic uncertainty:
  - background shape
  - neutral eff
  - tracking






# Motivation:

- $A_{CP} \sim 2\%$  in SM
- New Physics ( $A_{CP}$  up to 30%)
- $\sin 2\beta$  if no new physics

see Doug Wright's talk

## $B \rightarrow \phi K^{(*)}, \phi \pi$



PRD 65 (2002) 051101

$\sim 22.7 \times 10^6$  B pairs

$$A_{CP}(B^\pm \rightarrow \phi K^\pm) = -0.05 \pm 0.20 \pm 0.03$$

$$A_{CP}(B^\pm \rightarrow \phi K^{*\pm}) = -0.43_{-0.30}^{+0.36} \pm 0.06$$

$$A_{CP}(B^0 \rightarrow \phi K^{*0}) = 0.00 \pm 0.27 \pm 0.03$$

$\sim 22.7 \times 10^6$  B pairs

PRL 87 151801 (2001)

$$Br(B^\pm \rightarrow \phi K^{*\pm}) = (9.7_{-3.4}^{+4.2} \pm 1.7) \times 10^{-6}$$

$$Br(B^0 \rightarrow \phi K^{*0}) = (8.6_{-2.4}^{+2.8} \pm 1.1) \times 10^{-6}$$


$\sim 60 \times 10^6$  B pairs


$$Br(B^\pm \rightarrow \phi K^\pm) = (9.2 \pm 1.0 \pm 0.8) \times 10^{-6}$$

$$Br(B^0 \rightarrow \phi K_S^0) = (8.7_{-1.5}^{+1.7} \pm 0.9) \times 10^{-6}$$

$$Br(B^\pm \rightarrow \phi \pi^\pm) < 0.56 \times 10^{-6} \text{ (90\% C.L.)}$$

preliminary





- $A_{CP}$  – search for direct CPV
- access to  $\gamma$

$$A(B^+ \rightarrow \bar{D}^0 K^+) = |\bar{A}| e^{i\delta_1} (V_{cb}^* V_{us})$$

$$A(B^+ \rightarrow D^0 K^+) = |A| e^{i\delta_2} e^{i\gamma} (V_{ub}^* V_{cs})$$

$D^0$  reconstructed:

- in  $K\pi$ ,  $K\pi\pi^0$ ,  $K\pi\pi\pi$

- in  $K^+K^-$  (CP=+1 eigenstate)

$$R \equiv \frac{Br(B^- \rightarrow D^0 K^-)}{Br(B^- \rightarrow D^0 \pi^-)} = (8.31 \pm 0.35 \pm 0.20)\%$$

$$A_{CP} \equiv \frac{Br(B^- \rightarrow D_{CP}^0 K^-) - Br(B^+ \rightarrow D_{CP}^0 K^+)}{Br(B^- \rightarrow D_{CP}^0 K^-) + Br(B^+ \rightarrow D_{CP}^0 K^+)} = 0.17 \pm 0.23^{+0.09}_{-0.07}$$



$$B \rightarrow K^* \gamma$$

$\sim 22.7 \times 10^6$  B pairs

Phys Rev Lett 88 101805 (2002)



- $A_{CP} \leq 1\%$  in SM.
- SUSY can enhance to  $\sim 20\%$

Reconstruct

$$K^* \rightarrow K^+ \pi^-, K^+ \pi^0, K_s \pi^0, K_s \pi^+$$

$$Br(B^0 \rightarrow K^* \gamma) = (4.23 \pm 0.40 \pm 0.22) \times 10^{-5}$$

$$Br(B^+ \rightarrow K^{*+} \gamma) = (3.83 \pm 0.62 \pm 0.22) \times 10^{-5}$$




$$A_{CP}(B \rightarrow K^* \gamma) = -0.044 \pm 0.076 \pm 0.012$$

$$[-0.170, 0.082] \text{ (90\% C.L.)}$$



Other results: see talk on  $\alpha$

$\sim 88 \times 10^6$  B pairs



$$A_{CP}^{\rho K} = 0.19 \pm 0.14 \pm 0.11$$

$$A_{CP}^{\rho\pi} = -0.22 \pm 0.08 \pm 0.07$$



$$C_{\rho\pi} = 0.45^{+0.18}_{-0.19} \pm 0.09$$

$$S_{\rho\pi} = 0.16 \pm 0.25 \pm 0.07$$

$$\Delta C_{\rho\pi} = 0.38^{+0.19}_{-0.20} \pm 0.11$$

$$\Delta S_{\rho\pi} = 0.15 \pm 0.25 \pm 0.05$$

Factorization predicts 0.4



$\alpha$  via isospin analysis and direct CPV from time evolution of  $B \rightarrow \pi^+ \pi^-$

$$C_{\pi\pi} = -0.30 \pm 0.25 \pm 0.04$$

$$S_{\pi\pi} = 0.02 \pm 0.34 \pm 0.05$$

$$Br(B^0 \rightarrow \pi^+ \pi^-) = (4.7 \pm 0.6 \pm 0.2) \times 10^{-6}$$

$$Br(B^+ \rightarrow \pi^+ \pi^0) = (5.5 \pm 1.0 \pm 0.6) \times 10^{-6}$$

$$Br(B^0 \rightarrow \pi^0 \pi^0) < 3.6 \times 10^{-6}$$

Preliminary



# Summary and outlook

- First CP results from  $B^0 \rightarrow \rho\pi$
- Branching fractions and  $A_{CP}$  measured in many decays
- $A_{CP}$  precision achieved 5~20%
- No direct CPV signal yet ...
- watch for future updates

