

FIRST OSCILLATION RESULTS FROM THE NOVA EXPERIMENT




European Research Council
Established by the European Commission




Bruno Zamorano
Liverpool - 3rd February 2016



Why study neutrino oscillations?



- Second most abundant particle in the Universe and yet the worst understood
- Dark Matter aside, the only measured confirmation of Physics beyond the Standard Model
- ~20 000 neutrino papers since the discovery of neutrino oscillations
- Nobel prize 2015 and Breakthrough prize 2016
- Many open questions: CP violation (matter-antimatter asymmetry), mass ordering and mass scale, Dirac or Majorana...
- Oscillation parameters are, to our best knowledge, fundamental constants of Nature



Neutrino oscillations overview

PMNS
matrix

$$\begin{pmatrix} \nu_e \\ \nu_\mu \\ \nu_\tau \end{pmatrix} = \begin{pmatrix} U_{e1} & U_{e2} & U_{e3} \\ U_{\mu 1} & U_{\mu 2} & U_{\mu 3} \\ U_{\tau 1} & U_{\tau 2} & U_{\tau 3} \end{pmatrix} \begin{pmatrix} \nu_1 \\ \nu_2 \\ \nu_3 \end{pmatrix}$$

$$P_{\mu e} = \sum_{j,k} U_{ej}^* U_{\mu j} U_{\mu k}^* U_{ek} \exp\left(-i \frac{\Delta m_{jk}^2 L}{2E}\right)$$

Oscillations

Neutrino oscillations overview

PMNS matrix

$$\begin{pmatrix} \nu_e \\ \nu_\mu \\ \nu_\tau \end{pmatrix} = \begin{pmatrix} U_{e1} & U_{e2} & U_{e3} \\ U_{\mu 1} & U_{\mu 2} & U_{\mu 3} \\ U_{\tau 1} & U_{\tau 2} & U_{\tau 3} \end{pmatrix} \begin{pmatrix} \nu_1 \\ \nu_2 \\ \nu_3 \end{pmatrix}$$

$$P_{\mu e} = \sum_{j,k} U_{ej}^* U_{\mu j} U_{\mu k}^* U_{ek} \exp\left(-i \frac{\Delta m_{jk}^2 L}{2E}\right)$$


Oscillations

Atmospheric

Reactor

Solar

$$\begin{pmatrix} U_{e1} & U_{e2} & U_{e3} \\ U_{\mu 1} & U_{\mu 2} & U_{\mu 3} \\ U_{\tau 1} & U_{\tau 2} & U_{\tau 3} \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & c_{23} & s_{23} \\ 0 & -s_{23} & c_{23} \end{pmatrix} \times \begin{pmatrix} c_{13} & 0 & s_{13}e^{-i\delta} \\ 0 & 1 & 0 \\ -s_{13}e^{-i\delta} & 0 & c_{13} \end{pmatrix} \times \begin{pmatrix} c_{12} & s_{12} & 0 \\ -s_{12} & c_{12} & 0 \\ 0 & 0 & 1 \end{pmatrix}$$



$$\Delta m_{32}^2 \simeq \Delta m_{31}^2 \xrightarrow{Osc.max} L/E \approx 500 \text{ km/GeV}$$

$$\Delta m_{21}^2 = m_2^2 - m_1^2 \xrightarrow{Osc.max} L/E \approx 15000 \text{ km/GeV}$$

Importance of reactor result

$$\times \begin{pmatrix} c_{13} & 0 & s_{13}e^{-i\delta} \\ 0 & 1 & 0 \\ -s_{13}e^{-i\delta} & 0 & c_{13} \end{pmatrix} \times$$

θ_{13} : from unknown to best measured

$$\sin^2 \theta_{13} = 0.0219 \pm 0.0012$$

$$\theta_{13} \sim 8.5^\circ$$

CP violation $\iff \theta_{13} \neq 0$

A new door to probing CP violation, the mass ordering and the octant of θ_{23}

Importance of reactor result

$$\times \begin{pmatrix} c_{13} & 0 & s_{13}e^{-i\delta} \\ 0 & 1 & 0 \\ -s_{13}e^{-i\delta} & 0 & c_{13} \end{pmatrix} \times$$

CP violation $\iff \theta_{13} \neq 0$

θ_{13} : from unknown to best measured

$$\sin^2 \theta_{13} = 0.0219 \pm 0.0012$$

$$\theta_{13} \sim 8.5^\circ$$

A new door to probing CP violation, the mass ordering and the octant of θ_{23}

ν_e appearance:


$$P(\nu_\mu \rightarrow \nu_e) \simeq \sin^2(\theta_{23}) \sin^2(2\theta_{13}) \left(\Delta m_{32}^2 L / 4E \right) + \dots$$

Daya Bay reactor experiment:


$$\sin^2(2\theta_{13}) = 0.084 \pm 0.005$$

CP -violation? and matter effect modifications

θ_{23} octant



Mass hierarchy



Window to measure appearance: CP violation and mass-hierarchy

SUMMARY

The NOvA
experiment

Muon neutrino
disappearance

Electron neutrino
appearance

Future
sensitivities



The NOvA experiment




About 150 physicists from 6 countries (list growing...)
2 UK institutions: Sussex (8 people, since 2012) and UCL (6 people, just joined)
New collaborators and institutions welcome!




NuMI beam




- Capable of running at 700 kW
- Reliably running at ~500 kW
- World record 521 kW for 1 hour
- Full power by mid 2016
- 6.4×10^{20} POT delivered




- NuMI Off-Axis ν_e Appearance, the leading neutrino oscillation experiment in the NuMI beam



- NuMI Off-Axis ν_e Appearance, the leading neutrino oscillation experiment in the NuMI beam
- Two highly active scintillator detectors:
 - Far Detector: 14 kT, on surface
 - Near Detector: 300 T, 105 m underground



- NuMI Off-Axis ν_e Appearance, the leading neutrino oscillation experiment in the NuMI beam
- Two highly active scintillator detectors:
 - Far Detector: 14 kT, on surface
 - Near Detector: 300 T, 105 m underground
- 14 mrad off-axis narrowly peaked muon neutrino flux at 2 GeV, $L/E \sim 405 \text{ km/GeV}$
- ν_μ disappearance channel: $\theta_{23}, \Delta m^2_{32}$
- ν_e appearance channel: mass hierarchy, $\delta_{CP}, \theta_{13}, \theta_{23}$ and octant degeneracy



Also: neutrino cross sections at the ND, sterile neutrinos, supernovae...

Far detector dataset

- During the construction era, we **began collecting Physics data** with each far detector “diblock” (64 detector planes) as soon as it was **fully commissioned** and Physics-ready
- Thus, the FD size is **not static** throughout our data set


Protons-on-target in data set: 3.45×10^{20} POT

Fraction of detector operational: 79.4% (POT-weighted average)

Full-detector-equivalent exposure: 2.74×10^{20} POT-equiv




Partial Far Detector during construction
(6 diblock example)




Full Far Detector
(14 diblocks)

- Superb granularity for a detector this scale
- Outstanding event identification capability


NOvA Monte Carlo




1 radiation length = 38 cm
(6 cell depths, 10 cell widths)




10 μs of readout during NuMI beam pulse




10 μ s of readout during NuMI beam pulse




Full 550 μ s readout (colours show charge)




Zoomed on the 10 μs beam spill window



Zoomed on the time slice




Zoomed on the time slice




Zoomed on the time slice

Disappearance analysis in a nutshell...

Identify contained ν_μ CC events in both detectors



Measure both energy spectra




Infer on the oscillation mechanism from differences between near and far energy spectra


Disappearance analysis in a nutshell...

arXiv 1601.05037

Identify contained ν_μ CC events in both detectors



Measure both energy spectra



Infer on the oscillation mechanism from differences between near and far energy spectra


- Containment
- PID
- NC rejection
- Cosmic rejection

- Calibration
- Energy scale
- Hadronic energy

- Extrapolation
- Far / near ratio
- Best fit
- Contours
- Systematics



Signal selection: ν_μ CC

- First: basic containment cuts: require a buffer of no activity around the event
- Muon ID: 4-variable k-nearest neighbours algorithm to identify muons
 - Track length
 - dE/dx along track
 - Scattering along track
 - Track-only plane fraction
- Keep events with muon-ID > 0.75





Cosmic rejection

- From beam timing: a factor 10^5 reduction
- From event topology: 10^7
- For the remaining, a boosted decision tree is trained on data without beam
 - Direction
 - Position and length
 - Energy and number of hits in event
- Keep events with $\text{BDT} > 0.535$
- Final rate is measured directly using out-of-time data






Energy estimation

- Muon track: **length** $\Rightarrow E_\mu$
- Hadronic system:
 Σ Total visible E $\Rightarrow E_{\text{had}}$
- Reconstructed neutrino energy is the sum of them: **$E_\nu = E_\mu + E_{\text{had}}$**
- Energy resolution at beam peak: $\sim 7\%$




Kinematic variables in the ND after all cuts
(Sample purity in ND = 98%)



Far detector prediction

- Estimate the underlying **true energy distribution** of selected ND events
- Multiply by expected **Far/Near ratio** and $\nu_\mu \rightarrow \nu_\mu$ oscillation probability
- Convert FD true energy distribution into predicted FD reconstructed energy distribution
- Systematic uncertainties assessed by varying all MC-based steps




Systematics

Most of the systematic uncertainties have relatively **little influence on the result**

Hadronic energy systematic is one with a noticeable effect

(although the impact is reduced by ND-to-FD prediction procedure)






Tune hadronic energy in ND simulation to achieve better agreement in E_ν and E_{had}

Take the full **size of this tuning** as a systematic uncertainty

- Hadronic energy (14%, equiv. to 3% in neutrino energy)
- Neutrino flux (NA49 + beam transportation model)
- Absolute, relative normalisation (1%, 2%)
- Neutrino interactions (GENIE / Intranuke model)



- NC and tau neutrinos (100% each)
- Multiple calibration and light level systs. (Hit energy, fibre attenuation, threshold effects)
- Oscillation parameter uncertainties (current world knowledge)

Some of the selected candidates
(zoomed views, not showing the whole detector)




In the absence of oscillation, the expectation is 200 events (2 beam bkg., 1.4 cosmic bkg.)

33 selected events in the FD



Clear observation of ν_μ disappearance

Oscillation parameters



Normal hierarchy

$$\Delta m_{32}^2 = 2.52^{+0.20}_{-0.18} \times 10^{-3} \text{ eV}^2$$


Inverted hierarchy

$$\Delta m_{32}^2 = -2.56^{+0.19}_{-0.19} \times 10^{-3} \text{ eV}^2$$


NOvA measurement already compelling with less than 8% of nominal exposure!

Appearance analysis in a nutshell...

Identify contained
 ν_e CC events in
both detectors



Use ND candidates
to predict beam
backgrounds in the
FD




Interpret any FD
excess over
predicted
backgrounds as ν_e
appearance


Appearance analysis in a nutshell...

arXiv 1601.05022

Identify contained
 ν_e CC events in
both detectors



Use ND candidates
to predict beam
backgrounds in the
FD



Interpret any FD
excess over
predicted
backgrounds as ν_e
appearance


- Containment
- PID
- NC rejection
- Cosmic rejection

- Extrapolation
- Far / near ratio
- Systematics

- Exclusions
- Significance



Signal pre-selection: ν_e CC

- First: basic containment cuts: require sufficient distance from the largest reconstructed shower to the edges
- Then, cuts applied to
 - Shower length
 - Number of hits in event
 - Calorimetric energy
- All three related to the “size” of the event (we know well the range of energies any appearing ν_e might have)



Cosmic rejection

- Cut events with large transverse momentum (*Rejects downward-going cosmic showers*)
- The ν_e selectors themselves are very efficient at rejecting cosmic background
- Achieve a 1 in 10^8 rejection
- Expected cosmic background; 0.06 events, measured directly using out-of-time data



ν_e CC event identification


Two independent selection algorithms with very different designs

LID: Likelihood identification


dE/dX likelihoods calculated for **longitudinal and transverse** slices of leading shower under multiple particle hypotheses

Likelihoods feed an artificial neural network along with **kinematic and topological info**: e.g., energy near vertex, shower angle, vertex-to-shower gap, ...

Color: p.d.f. for dE/dx in each plane (e^- assumption)



Likelihoods calculated for each red and yellow region




red: longitudinal slices
yellow: transverse slices

LEM: Library event matching

Spatial pattern of energy depositions is compared to that of 10^8 simulated events (“library”)

Key properties of the **best matched library event** (e.g., fraction that are signal events) are input into a decision tree to form discriminant

Left panels: candidate event, both views
Right panels: best-matched library event, both views
Middle panels: an intermediate step in calculating the match quality



- Identical performance as measured with efficiency, S/\sqrt{B} and sensitivity to oscillation parameters

Prior to unblinding, decided to **show both results and use LID as primary**

LID and LEM distributions for ND data and simulation



- Good agreement over full range

Total BKG	ν_e - CC (beam)	NC	ν_μ - CC	ν_τ - CC	Cosmic
0.94 ± 0.09	0.46	0.35	0.05	0.02	0.06
\pm few % depending on osc. param.					

Range of signal predictions


- NH, $\delta_{CP} = 3\pi/2$, $\theta_{23} = \pi/4 \rightarrow 5.6 \pm 0.7$
- IH, $\delta_{CP} = \pi/2$, $\theta_{23} = \pi/4 \rightarrow 2.2 \pm 0.3$

NOvA Preliminary



Far detector prediction

- **ND data** are translated to **FD background expectation** in each energy bin, using far/near ratios from simulation
- **FD signal expectation** is pinned to the ND-selected $\nu\mu$ - CC spectrum
- Most **systematics** are assessed **via variations** in the far/near ratios



Signal efficiency relative to containment cuts: 35%

Expected overlap in LID/LEM samples of 62%

(Expected differences in which events each technique selects)

After all selection, **0.7% of NC** events remain, relative to those after containment

Far detector selected events



LID: 6 ν_e candidates

3.3 σ significance for ν_e appearance

LEM: 11 ν_e candidates




5.5 σ significance for ν_e appearance

(All 6 LID events present in LEM set)













It feels me with pride showing our hall of fame in
Liverpool













(I swear this was not fabricated for this seminar)



Electron neutrino appearance




Electron neutrino appearance



Result using LID selector

For allowed regions of δ_{CP} and θ_{13}

- Feldman & Cousins procedure applied
- Solar oscillation parameters varied
- Δm^2_{32} varied by new NOvA result
- $\sin^2\theta_{23} = 0.5$




Result using LID selector

Applying global reactor constraint of



$$\sin^2 \theta_{13} = 0.0219 \pm 0.0012$$

- Again, apply Feldman-Cousins to interpret $-2\Delta\ln L$
- Converted into significance (*steps due to discrete nature of counting experiment*)
- Using ν_μ constraint for $\sin^2 \theta_{23}$



Inverted hierarchy for $\delta_{CP} \in [0, 0.9\pi]$ is mildly disfavoured ($> 1\sigma$)


Result using LEM selector



- Inverted hierarchy is disfavoured at 2.2σ for all δ_{CP}
- NH for $\delta_{CP} \in [0, 1.3\pi]$ is mildly disfavoured ($> 1\sigma$)


LID/LEM consistency

- Both prefer normal hierarchy
- Both prefer δ_{CP} near $3\pi/2$
- Given the expected correlations, the observed event counts yield a reasonable **mutual p-Value of 10%**





The specific point IH, $\delta_{CP} = \pi/2$ is disfavoured at 1.6σ (LID), 3.2σ (LEM)

- Potential to exclude maximal mixing, depending on Nature's choice
- Leading measurement in both Δm^2_{32} and $\sin^2\theta_{23}$ for nominal sensitivity
- Measurements in the anti-neutrino channel: CPT tests





COMBINING MUON AND ELECTRON NEUTRINO ANALYSES



Best case scenario: NOvA simultaneously measures the mass ordering, CP violation and octant information!


COMBINING MUON AND ELECTRON NEUTRINO ANALYSES




Degenerate case: mass ordering and CP violation are coupled, but the octant information is not

MASS HIERARCHY AND CP-VIOLATION

NOvA hierarchy resolution

 $\sin^2 2\theta_{13}=0.095, \sin^2 2\theta_{23}=1.00$
 36×10^{20} POT




NOvA CPV determination

 $\sin^2 2\theta_{13}=0.095, \sin^2 2\theta_{23}=1.00$
 36×10^{20} POT


3+3 years (ν_μ + anti- ν_μ): 2 sigma in
about 30% of the δ_{CP} range

Only 1.5 sigma in 10% of the range

COMBINATION WITH T2K



Combining with T2K: At least 1 sigma for the whole δ_{CP} range

With T2K: 1.5 sigma in 25%

- Far and near detector completed in August 2014
- The NuMI beam intensity is steadily increasing, setting new records on a weekly basis
- First analysis dataset included 7.8% of full exposure (40% of a standard running year)
- Other NOvA physics: programme underway
 - Neutrino cross-sections
 - Sterile neutrinos, non-standard interactions, CPT tests
 - Supernova neutrinos
 - Dark matter and monopole searches
 - *And more...*

- Far and near detector completed in August 2014
- The NuMI beam intensity is steadily increasing, setting new records on a weekly basis
- First analysis dataset included 7.8% of full exposure (40% of a standard running year)
- Other NOvA physics: programme underway
 - Neutrino cross-sections
 - Sterile neutrinos, non-standard interactions, CPT tests
 - Supernova neutrinos
 - Dark matter and monopole searches
 - *And more...*

With 2.74×10^{20} POT exposure...

- Unambiguous ν_μ disappearance signature
- 8% measurement of atmospheric mass splitting
- Consistent with maximal mixing
- ν_e appearance signal at 3.3σ for primary selector, 5.5σ for secondary selector
- At max mixing, disfavour IH for $\delta_{CP} \in [0.1\pi, 0.5\pi]$ at 90% with primary selector. Further preference for NH with secondary selector

Stay tuned!



THANK YOU FOR YOUR ATTENTION

Public live event display:
<http://nusoft.fnal.gov/nova/public/>



www-nova.fnal.gov