

# An Energy Recovery Linac for Energy-Frontier eh Scattering at CERN: the LHeC and the FCC-eh


Oliver Brüning (CERN), Max Klein (U Liverpool), Daniel Schulte (CERN)

for the LHeC/PERLE/FCCeh Collaboration




Talk at ICHEP Seoul, 6 July, 2018

# 60 GeV Electron ERL added to LHC



**Concurrent operation to pp, LHC/FCC become 3 beam facilities. Power limit: 100 MW**  
 $10^{34} \text{ cm}^{-2} \text{ s}^{-1}$  luminosity and factor of 15/120 (LHC/FCCeh) extension of  $Q^2$ ,  $1/x$  reach  
 1000 times HERA luminosity. It therefore extends up to  $x \sim 1$ .  
 Four orders of magnitude extension in deep inelastic lepton-nucleus (ion) scattering.



## CERN Referees

### **Ring Ring Design**

Kurt Huebner (CERN)  
Alexander N. Skrinsky (INP Novosibirsk)  
Ferdinand Willeke (BNL)

### **Linac Ring Design**

Reinhard Brinkmann (DESY)  
Andy Wolski (Cockcroft)  
Kaoru Yokoya (KEK)

### **Energy Recovery**

Georg Hoffstaetter (Cornell)  
Ilan Ben Zvi (BNL)

### **Magnets**

Neil Marks (Cockcroft)  
Martin Wilson (CERN)

### **Interaction Region**

Daniel Pitzl (DESY)  
Mike Sullivan (SLAC)

### **Detector Design**

Philippe Bloch (CERN)  
Roland Horisberger (PSI)

### **Installation and Infrastructure**

Sylvain Weisz (CERN)

### **New Physics at Large Scales**

Cristinel Diaconu (IN2P3 Marseille)  
Gian Giudice (CERN)

Michelangelo Mangano (CERN)

### **Precision QCD and Electroweak**

Guido Altarelli (Roma)  
Vladimir Chekelian (MPI Munich)  
Alan Martin (Durham)

### **Physics at High Parton Densities**

Alfred Mueller (Columbia)  
Raju Venugopalan (BNL)  
Michele Arneodo (INFN Torino)

Published 600 pages conceptual design report (CDR) written by 200 authors from 60 Institutes and refereed by 24 world experts on physics, accelerator and detector, which CERN had invited.

<b>7</b>	<b>Linac-Ring Collider</b>		<b>317</b>
7.1	Basic parameters and configurations		317
7.1.1	General considerations		317
7.1.2	ERL performance and layout		318
7.1.3	Polarisation	<b>Performance</b>	327
7.1.4	Pulsed linacs		329
7.1.5	Higher-energy LHeC ERL option		329
7.1.6	$\gamma$ - $p/A$ Option		330
7.1.7	Summary of basic parameters and configurations		331
7.2	Interaction region		332
7.2.1	Layout		333
7.2.2	Optics	<b>Interaction Region</b>	334
7.2.3	Modifications for $\gamma p$ or $\gamma A$		340
7.2.4	Synchrotron radiation and absorbers		340
7.3	Linac lattice and impedance		348
7.3.1	Overall layout		348
7.3.2	Linac layout and lattice	<b>Optics</b>	350
7.3.3	Beam break-up		356
7.3.4	Imperfections		368
7.3.5	Touschek scattering		369
7.4	Performance as a Linac-Ring electron-ion collider		369
7.4.1	Heavy nuclei, e-Pb collisions		370
7.4.2	Electron-deuteron collisions	<b>Ions</b>	370
7.5	Polarised-electron injector for the Linac-Ring LHeC		370
7.6	Spin Rotator		372
7.6.1	Introduction		372
7.6.2	LHeC spin rotator options		373
7.6.3	Polarimetry	<b>Polarisation (80-90%)</b>	376
7.6.4	Conclusions and Outlook		377
7.7	Positron options for the Linac-Ring LHeC		379
7.7.1	Motivation		379
7.7.2	LHeC Linac-Ring $e^+$ requirements	<b>Positrons</b>	380
7.7.3	Mitigation schemes		380
7.7.4	Cooling of positrons		381
7.7.5	Production schemes		382
7.7.6	Conclusions on positron options for the Linac-Ring LHeC		388

**CDR: VERY detailed design of the LHeC Linac (and Ring) – Ring Collider, + components, CE..**



J.L.Abelleira Fernandez<sup>16,23</sup>, C.Adolphsen<sup>57</sup>, P.Adzic<sup>74</sup>, A.N.Akay<sup>03</sup>, H.Aksakal<sup>39</sup>, J.L.Albacete<sup>52</sup>, B.Allanach<sup>73</sup>, S.Alekhin<sup>17,54</sup>, P.Allport<sup>24</sup>, V.Andreev<sup>34</sup>, R.B.Appleby<sup>14,30</sup>, E.Arikan<sup>39</sup>, N.Armento<sup>53,a</sup>, G.Azuelos<sup>33,64</sup>, M.Bai<sup>37</sup>, D.Barber<sup>14,17,24</sup>, J.Bartels<sup>18</sup>, O.Behnke<sup>17</sup>, J.Behr<sup>17</sup>, A.S.Belyaev<sup>15,56</sup>, I.Ben-Zvi<sup>37</sup>, N.Bernard<sup>25</sup>, S.Bertolucci<sup>16</sup>, S.Bettoni<sup>16</sup>, S.Biswal<sup>41</sup>, J.Blümlein<sup>17</sup>, H.Böttcher<sup>17</sup>, A.Bogacz<sup>36</sup>, C.Bracco<sup>16</sup>, J.Bracinik<sup>06</sup>, G.Brandt<sup>44</sup>, H.Braun<sup>65</sup>, S.Brodsky<sup>57,b</sup>, O.Brüning<sup>16</sup>, E.Bulyak<sup>12</sup>, A.Buniatyan<sup>17</sup>, H.Burkhardt<sup>16</sup>, I.T.Cakir<sup>02</sup>, O.Cakir<sup>01</sup>, R.Calaga<sup>16</sup>, A.Caldwell<sup>70</sup>, V.Cetinkaya<sup>01</sup>, V.Chekelian<sup>70</sup>, E.Ciapala<sup>16</sup>, R.Ciftci<sup>01</sup>, A.K.Ciftci<sup>01</sup>, B.A.Cole<sup>38</sup>, J.C.Collins<sup>48</sup>, O.Dadoun<sup>42</sup>, J.Dainton<sup>24</sup>, A.De.Roeck<sup>16</sup>, D.d'Enterria<sup>16</sup>, P.DiNezza<sup>72</sup>, M.D'Onofrio<sup>24</sup>, A.Dudarev<sup>16</sup>, A.Eide<sup>60</sup>, R.Enberg<sup>63</sup>, E.Eroglu<sup>62</sup>, K.J.Eskola<sup>21</sup>, L.Favart<sup>08</sup>, M.Fitterer<sup>16</sup>, S.Forte<sup>32</sup>, A.Gaddi<sup>16</sup>, P.Gambino<sup>59</sup>, H.García Morales<sup>16</sup>, T.Gehrmann<sup>69</sup>, P.Gladkikh<sup>12</sup>, C.Glasman<sup>28</sup>, A.Glazov<sup>17</sup>, R.Godbole<sup>35</sup>, B.Goddard<sup>16</sup>, T.Greenshaw<sup>24</sup>, A.Guffanti<sup>13</sup>, V.Guzey<sup>19,36</sup>, C.Gwenlan<sup>44</sup>, T.Han<sup>50</sup>, Y.Hao<sup>37</sup>, F.Haug<sup>16</sup>, W.Herr<sup>16</sup>, A.Hervé<sup>27</sup>, B.J.Holzer<sup>16</sup>, M.Ishitsuka<sup>58</sup>, M.Jacquet<sup>42</sup>, B.Jeaneret<sup>16</sup>, E.Jensen<sup>16</sup>, J.M.Jimenez<sup>16</sup>, J.M.Jowett<sup>16</sup>, H.Jung<sup>17</sup>, H.Karadeniz<sup>02</sup>, D.Kayran<sup>37</sup>, A.Kilic<sup>62</sup>, K.Kimura<sup>58</sup>, R.Klees<sup>75</sup>, M.Klein<sup>24</sup>, U.Klein<sup>24</sup>, T.Kluge<sup>24</sup>, F.Kocak<sup>62</sup>, M.Korostelev<sup>24</sup>, A.Kosmicki<sup>16</sup>, P.Kostka<sup>17</sup>, H.Kowalski<sup>17</sup>, M.Kraemer<sup>75</sup>, G.Kramer<sup>18</sup>, D.Kuchler<sup>16</sup>, M.Kuze<sup>58</sup>, T.Lappi<sup>21,c</sup>, P.Laycock<sup>24</sup>, E.Levichev<sup>40</sup>, S.Levonian<sup>17</sup>, V.N.Litvinenko<sup>37</sup>, A.Lombardi<sup>16</sup>, J.Maeda<sup>58</sup>, C.Marquet<sup>16</sup>, B.Mellado<sup>27</sup>, K.H.Mess<sup>16</sup>, A.Milanese<sup>16</sup>, J.G.Milhano<sup>76</sup>, S.Moch<sup>17</sup>, I.I.Morozov<sup>40</sup>, Y.Muttoni<sup>16</sup>, S.Myers<sup>16</sup>, S.Nandi<sup>55</sup>, Z.Nergiz<sup>39</sup>, P.R.Newman<sup>06</sup>, T.Omori<sup>61</sup>, J.Osborne<sup>16</sup>, E.Paoloni<sup>49</sup>, Y.Papaphilippou<sup>16</sup>, C.Pascaud<sup>42</sup>, H.Paukkunen<sup>53</sup>, E.Perez<sup>16</sup>, T.Pieloni<sup>23</sup>, E.Pilicer<sup>62</sup>, B.Pire<sup>45</sup>, R.Placakyte<sup>17</sup>, A.Polini<sup>07</sup>, V.Ptitsyn<sup>37</sup>, Y.Pupkov<sup>40</sup>, V.Radescu<sup>17</sup>, S.Raychaudhuri<sup>35</sup>, L.Rinolfi<sup>16</sup>, E.Rizvi<sup>71</sup>, R.Rohini<sup>35</sup>, J.Rojo<sup>16,31</sup>, S.Russenschuck<sup>16</sup>, M.Sahin<sup>03</sup>, C.A.Salgado<sup>53,a</sup>, K.Sampej<sup>58</sup>, R.Sassot<sup>09</sup>, E.Sauvan<sup>04</sup>, M.Schaefer<sup>75</sup>, U.Schneekloth<sup>17</sup>, T.Schörner-Sadenius<sup>17</sup>, D.Schulte<sup>16</sup>, A.Senol<sup>22</sup>, A.Seryi<sup>44</sup>, P.Sievers<sup>16</sup>, A.N.Skrinsky<sup>40</sup>, W.Smith<sup>27</sup>, D.South<sup>17</sup>, H.Spiesberger<sup>29</sup>, A.M.Stasto<sup>48,d</sup>, M.Strikman<sup>48</sup>, M.Sullivan<sup>57</sup>, S.Sultansoy<sup>03,e</sup>, Y.P.Sun<sup>57</sup>, B.Surrow<sup>11</sup>, L.Szymanowski<sup>66,f</sup>, P.Taels<sup>05</sup>, I.Tapan<sup>62</sup>, T.Tasci<sup>22</sup>, E.Tassi<sup>10</sup>, H.Ten.Kate<sup>16</sup>, J.Terron<sup>28</sup>, H.Thiesen<sup>16</sup>, L.Thompson<sup>14,30</sup>, P.Thompson<sup>06</sup>, K.Tokushuku<sup>61</sup>, R.Tomás García<sup>16</sup>, D.Tommasini<sup>16</sup>, D.Trbojevic<sup>37</sup>, N.Tsoupas<sup>37</sup>, J.Tuckmantel<sup>16</sup>, S.Turkoz<sup>01</sup>, T.N.Trinh<sup>47</sup>, K.Tywoniuk<sup>26</sup>, G.Unel<sup>20</sup>, T.Ullrich<sup>37</sup>, J.Urakawa<sup>61</sup>, P.VanMechelen<sup>05</sup>, A.Variola<sup>52</sup>, R.Veness<sup>16</sup>, A.Vivoli<sup>16</sup>, P.Vobly<sup>40</sup>, J.Wagner<sup>66</sup>, R.Wallny<sup>68</sup>, S.Wallon<sup>43,46,f</sup>, G.Watt<sup>69</sup>, C.Weiss<sup>36</sup>, U.A.Wiedemann<sup>16</sup>, U.Wienands<sup>57</sup>, F.Willeke<sup>37</sup>, B.-W.Xiao<sup>48</sup>, V.Yakimenko<sup>37</sup>, A.F.Zarnecki<sup>67</sup>, Z.Zhang<sup>42</sup>, F.Zimmermann<sup>16</sup>, R.Zlebcik<sup>51</sup>, F.Zomer<sup>42</sup>

LHeC Study group and CDR authors 2014

“you never walk alone”



## A Higgs Facility Resolving the Substructure of Matter

Update on the 2012 LHeC Report  
on the Physics and Design Concepts for Machine and Detector

---

LHeC Collaboration



Submitted to J.Phys. G

**DRAFT**

## New LHeC Document

Reasons: CERN Mandate and


- Higgs: higher luminosity goal
- LHC: no BSM + precision
- New ep/eA Physics prospects
- eh appeared with FCC design
- Updates on IR, CE, ...
- Insight from FCC study
- Vision for eh with HE LHC
- Options for ERL based Physics beyond eh scattering.
- Low energy test facility PERLE
- Implementation of ERL at CERN

Preparation for early 2019

## Submissions of eh Papers to European Strategy:

- FCC eh as part of FCC hh
- LHeC with HL
- PERLE technology development

# Physics with Energy Frontier DIS



## Raison(s) d'etre of the LHeC

Cleanest High Resolution  
Microscope: QCD Discovery

Empowering the LHC  
Search Programme

Transformation of LHC into  
high precision Higgs facility


Discovery (top, H, heavy  $\nu$ 's..)  
Beyond the Standard Model

A Unique  
Nuclear Physics Facility



# Location, Footprint, Use of the Electron Racetrack

e beam external to LHC. Location suitable for both HL and HE LHC.




- $U(\text{ERL}) = 1/n U(\text{LHC})$ : 60 GeV: 1/3
- BSM, top, Higgs, Low x all want maximum  $E_e$
- Cost goes almost linearly down with  $E_e$

For FCC can realise ep/A collisions  
With IR at point L, not far from CERN  
 $U(\text{ERL}) = 1/11 U(\text{FCC})$

**Energy – Cost – Physics – Footprint** are being reinvestigated for EU strategy

# Civil Engineering for ERL



For LHC: re-use IP2. For FCC: deeper shafts, new IP cavern  
 Refinement of CE study, see J Osborne Amsterdam FCC week  
 and see Matt Stuart at LHeC Workshop, 28.6.18 at Orsay

For matched electron and proton beam sizes:

$$\mathcal{L} \propto \frac{N_p}{\epsilon_p} \frac{1}{\beta_p} I_e H_{geom} H_{b-b} H_{coll}$$

The diagram illustrates the luminosity formula  $\mathcal{L} \propto \frac{N_p}{\epsilon_p} \frac{1}{\beta_p} I_e H_{geom} H_{b-b} H_{coll}$  with the following annotations:

- $N_p$  (red oval) is linked to **Proton beam brilliance** (red box).
- $\epsilon_p$  (red oval) is linked to **Proton beam brilliance** (red box).
- $\beta_p$  (orange oval) is linked to **Proton ring design** (orange box).
- $I_e$  (green oval) is linked to **Electron beam current** (green box).
- $H_{geom}$  (brown oval) is linked to **Hourglass effect** (brown box).
- $H_{b-b}$  (blue oval) is linked to **Beam-beam effect** (blue box).
- $H_{coll}$  (purple oval) is linked to **Fill pattern matching** (purple box).

→  **$O(1ab^{-1})$  in a decade of operation**, see D Schulte, FCC week at Rome, 2016



# Luminosity for LHeC, HE-LHeC and FCC-ep

parameter [unit]	LHeC CDR	ep at HL-LHC	ep at HE-LHC	FCC-he
$E_p$ [TeV]	7	7	12.5	50
$E_e$ [GeV]	60	60	60	60
$\sqrt{s}$ [TeV]	1.3	1.3	1.7	3.5
bunch spacing [ns]	25	25	25	25
protons per bunch [ $10^{11}$ ]	1.7	2.2	2.5	1
$\gamma\epsilon_p$ [ $\mu\text{m}$ ]	3.7	2	2.5	2.2
electrons per bunch [ $10^9$ ]	1	2.3	3.0	3.0
electron current [mA]	6.4	15	20	20
IP beta function $\beta_p^*$ [cm]	10	7	10	15
hourglass factor $H_{geom}$	0.9	0.9	0.9	0.9
pinch factor $H_{b-b}$	1.3	1.3	1.3	1.3
proton filling $H_{coll}$	0.8	0.8	0.8	0.8
luminosity [ $10^{33}\text{cm}^{-2}\text{s}^{-1}$ ]	1	8	12	15


Oliver Brüning<sup>1</sup>, John Jowett<sup>1</sup>, Max Klein<sup>2</sup>,

Dario Pellegrini<sup>1</sup>, Daniel Schulte<sup>1</sup>, Frank Zimmermann<sup>1</sup>

EDMS 17979910 | FCC-ACC-RPT-0012

Contains update on eA:  
6  $10^{32}$  in e-Pb for LHeC.

# Collider Luminosities vs Year (pp and ep)



# FCC-eh ERL Configuration:

[Daniel Schulte]

## Performance Simulations for FCC-ep:

Parameter	Unit	Protons	Electrons
Beam energy	GeV	50000	60
Normalised emittance	$\mu\text{m}$	2.2 $\rightarrow$ 1.1	10
IP betafunction	mm	150	42 $\rightarrow$ 52
Nominal RMS beam size	$\mu\text{m}$	2.5 $\rightarrow$ 1.8	1.9 $\rightarrow$ 2.1
Waist shift	mm	0	65 $\rightarrow$ 70
Bunch population	$10^{10}$	10 $\rightarrow$ 5	0.31
Bunch spacing	ns	25	25
Luminosity	$10^{33}\text{cm}^{-2}\text{s}^{-1}$	18.3 $\rightarrow$ 14.3	
Int. luminosity per 10 years	$[\text{ab}^{-1}]$	1.2	

# DA and Status of Lattices

## LHeC


- Based on HL-LHC lattice (round optics  $B^*=15$  cm in IR1 and IR5)
- New low- $B^*$  IR (IR2)
- ATS-scheme implemented in 3 low- $B^*$  IRs
- Previous DA studies were implemented for different IR options
- Update: Studies with errors in IR1/IR5 magnets and new magnet design for IR2

## FCC-eh

- Based on FCC-hh lattice ( $B^*=30$  cm in IRA and IRG)
- New low- $B^*$  IR (IRL)
- No ATS-scheme
- Extensive DA studies have been performed for FCC-hh lattice
- Update: Implement same techniques for FCC-eh


Update of Lattice and IR Design Study: see E Cruz, R Martin and B Parker at Orsay Workshop

# Recent Progress on IR Magnets




- The yoke and a small fine tuning coil add 5% to the bare coil gradient and create zero field sweet spot region just outside coil structure.
- If this were made an actively shielded coil the gradient would have instead dropped by 7% and the “septum region” made 10% thicker.
- A weak fine tuning coil allows to adjust field in the slot and compensate for saturation effects.

- Self-supporting coil structure presents smallest possible “septum” region between the beams.
- Quadrupole symmetric yoke with deep slot cut out regions bypasses magnetic flux around the near zero field space used for beam separation.
- Yoke slots are sized to cleanly pass the synrad fan cleanly to far from IP and thereby avoid adding backscatter background.






# LHeC-FEL – world's best X-ray source



peak brilliance, LHeC-FEL compared with state-of-the-art






## SASE FEL simulation results

evolution of the pulse power along the undulator

spatial (temporal) profile of the radiation pulse


wavelength spectrum of the radiation



for  $\lambda=0.45 \text{ \AA}$  ( $K = 4.24$ )

Z. Nergiz, F. Zimmermann, H. Aksakal

average brilliance, LHeC-FEL compared with state-of-the-art






**OPEN ACCESS**

**IOP** Publishing

Journal of Physics G: Nuclear and Particle Physics


J. Phys. G: Nucl. Part. Phys. **00** (2018) 000000 (71 pp)

# PERLE. Powerful energy recovery linac for experiments. Conceptual design report

D Angal-Kalinin<sup>1</sup>, G Arduini<sup>2</sup>, B Auchmann<sup>2</sup>, J Bernauer<sup>3</sup>,  
A Bogacz<sup>4</sup>, F Bordry<sup>2</sup>, S Bousson<sup>5</sup>, C Bracco<sup>2</sup>, O Brüning<sup>2</sup>,  
R Calaga<sup>2</sup>, K Cassou<sup>6</sup>, V Chetvertkova<sup>2</sup>, E Cormier<sup>7</sup>, E Daly<sup>4</sup>,  
D Douglas<sup>4</sup>, K Dupraz<sup>6</sup> , B Goddard<sup>2</sup>, J Henry<sup>4</sup>, A Hutton<sup>4</sup>,  
E Jensen<sup>2</sup>, W Kaabi<sup>6</sup>, M Klein<sup>8</sup> , P Kostka<sup>8</sup>, N Lasheras<sup>2</sup>,  
E Levichev<sup>9</sup>, F Marhauser<sup>4</sup>, A Martens<sup>6</sup>, A Milanese<sup>2</sup>,  
B Militsyn<sup>1</sup>, Y Peinaud<sup>6</sup>, D Pellegrini<sup>2</sup>, N Pietralla<sup>10</sup>,  
Y Pupkov<sup>9</sup>, R Rimmer<sup>4</sup>, K Schirm<sup>2</sup>, D Schulte<sup>2</sup>, S Smith<sup>1</sup>,  
A Stocchi<sup>6</sup>, A Valloni<sup>2</sup>, C Welsch<sup>8</sup>, G Willering<sup>2</sup>, D Wollmann<sup>2</sup>,  
F Zimmermann<sup>2</sup> and F Zomer<sup>6</sup>

# Powerful ERL for Experiments at Orsay

- 2 Linacs (Four 5-Cell 801.58 MHz SC cavities)
- 3 turns (160 MeV/turn)
- Max. beam energy 500 MeV at 20mA  $\rightarrow$  10 MW



cf Walid Kaabi at Amsterdam FCC

**New SCRF, High Intensity (100 x ELI) ERL Development Facility with unique low E Physics**



# Why PERLE [as seen from LHeC]?



## FUNDAMENTAL MOTIVATION:

- **Validation of key LHeC Design Choices**
- **Build up expertise in the design and operation for a facility with a fundamentally new operation mode:**
  - ERLs are circular machines with tolerances and timing requirements similar to linear accelerators (no 'automatic' longitudinal phase stability, etc.)
- **Proof validity of fundamental design choices:**
  - Multi-turn recirculation (other existing ERLs have only 1-2 passages)
  - Implications of high current operation ( $2 * 3 * [6\text{mA} - 25\text{mA}] \rightarrow 30\text{-}150\text{mA}!!$ )
- **Verify and test machine and operation tolerances before designing a large scale facility**
  - Tolerances in terms of field quality of the arc magnets and cavity alignment
  - Required RF phase stability (RF power) and LLRF requirements
  - Halo and beam loss tolerances

# Frequency Choice






Cost, dynamic heat losses, resistance,  $Q_0$ ... point to  $f < 1$  GHz (F Marhauser, Orsay 2/17)

Beam beam interactions unstable for  $f > 1$  GHz (D Schulte, D Pellegrini March 2013)

Compatibility with LHC: **Decision for 802 MHz** ( E Jensen CI Workshop 1/2015, FM input)


# Towards PERLE: 802 MHz cavity, Source, Cryomodule, Magnets

First 802 MHz cavity successfully built (Jlab)



BINP, CERN, Daresbury/Liverpool, Jlab, Orsay, +  
CDR 1705.08783 [J.Phys G] → TDR in 2019







# Summary and Plans

- Following the CDR and the Higgs Boson Discovery in 2012, Physics from LHC, as well as Technology Developments, the ERL Concept for the LHeC has been updated to design for  $10^{34} \text{ cm}^{-2} \text{ s}^{-1}$  ep scattering with the HL LHC, HE LHC and the FCCeh.
- New physics (Higgs, BSM, top) studies have substantially widened the scope.
- The LHeC is designed for concurrent ep operation, with pp, in order to accumulate  $O(1) \text{ ab}^{-1}$  of integrated luminosity, at a power limit of 100 MW.
- New studies on lattice and the IR optics and configuration support this as a realistic goal.
- The heart of ep with HL LHC, HE LHC and FCC-eh is an energy recovery electron linac.
- A first SC cavity of 802 MHz frequency has been designed and successfully tested showing stability of up to 29 MV/m and a weak  $Q_0$ -gradient dependence around  $3 \cdot 10^{10}$ , exceeding the design goal. Single cells will be infused with N, + the CM designed
- A PERLE CDR has been published in 2017 and a TDR is scheduled for 2019. PERLE will be the first 10 MW, multi-turn 802 MHz facility and is suited to accompany the development of the LHeC as a technology development facility, with low E physics.
- PERLE and LHeC (HL/HE) inputs are under preparation for the HEP strategy update + beyond

**Most up-to-date Information:** <https://indico.cern.ch/event/698368/>

Workshop: LHeC/FCCeh and PERLE  
End of June at Orsay near Paris



# Electrons for the LHC

## LHeC/FCCeh and PERLE Workshop

June 27-29, 2018  
LAL-Orsay, France

**Organising Committee:**  
Nestor Armesto (USC)  
Oliver Brüning (CERN)  
Walid Kaabi (LAL)  
Uta Klein (Liverpool)  
Zhiqing Zhang (LAL)

**Advisory Committee:**  
Sergio Bertolucci (Bologna)  
Nicola Bianchi (INFN)  
Frederick Bondur (CERN)  
Oliver Brüning (CERN)  
Stanley Brodzsky (SLAC)  
Hsiang Chen (IHEP Beijing)  
Eckhard Eisen (CERN)  
Siddhant Fome (Milano)  
Andrew Horton (Jefferson Lab)

**Coordination Group:**  
Nestor Armesto (Santiago de Compostela)  
Gianluigi Arduini (CERN)  
Oliver Brüning (CERN)  
Andrea Gaddi (CERN)  
Erik Jensen (CERN)  
Bruce Mellado (Wits)  
Paul Newman (Birmingham)  
Daniel Schulte (CERN)  
Frank Zimmermann (CERN)

**Physics Convenors:**  
Daniel Beresford (Santiago de Compostela)  
Gergely Ásványi (München)  
Gert Baur (CERN)  
Gerrit Bevilacqua (CERN)  
Oliver Brüning (CERN)  
Giovanni Casella (CERN)  
Giovanni De Simone (CERN)  
Eliana Eidelson (CERN)  
Paul Newman (Birmingham)  
Sergio Bertolucci (Bologna)  
Siddhant Fome (Milano)  
Zhiqing Zhang (LAL)

<https://indico.cern.ch/event/698368/>





**New and Updates on**  
**Physics (PDFs, QCD, H, t, BSM, eA + Relation eh-hh..)**  
**Accelerator: IR, Optics, Lattice, Cost-Energy, CE..**  
**Detector: the GPD and its fwd and bwd detectors**  
**PERLE: Source, Injector, Cavity, Cryomodule,.. Physics**  
**Project Development towards the ES2020:**  
**LHeC + FCCeh+ PERLE input 12/18. PERLE TDR in 2019.**

<http://lhec.web.cern.ch>

backup

# Determination of SM Higgs Couplings, **HL-LHC** and **LHeC** → **LHC**

$\delta\kappa/\kappa$  [%]



- LHC
- LHeC
- ep+pp
- ep+pp, no thy unc

J. De Blas, M.+U. Klein, 16.4.2018

LHC: ATLAS prospects PUB Note 2014-016

ttH at LHeC to 15%

**The addition of ep to pp (LHeC to LHC (HL,HE) and FCC-eh to FCC-pp) transforms these machines into precision Higgs facilities. Vital complementarity with  $e^+e^-$  (JdB Amsterdam)**

Note that the HL LHC prospects are being updated (HL/HE LHC Physics workshop).



# Large Hadron Electron Collider on one page

$E_e = 10\text{-}60\text{ GeV}$ ,  $E_p = 1\text{-}7\text{ TeV}$ :  $\sqrt{s} = 200 - 1300\text{ GeV}$ . **Kinematics**:  $0 < Q^2 < s$ ,  $1 > x \geq 10^{-6}$  (DIS)  
Electron Polarisation  $P = \pm 80\%$ . Positrons: significantly lower intensity, unpolarised  
**Luminosity**:  $O(10^{34})\text{ cm}^{-2}\text{ s}^{-1}$ . integrated  $O(1)\text{ ab}^{-1}$  for HL LHC and  $2\text{ ab}^{-1}$  for HE LHC/FCCeh  
e-ions  $6 \cdot 10^{32}\text{ cm}^{-2}\text{ s}^{-1}$   $O(10)\text{ fb}^{-1}$  in ePb .  $O(1)\text{ fb}^{-1}$  for ep  $F_L$  measurements



**Physics**: QCD: develop+break? The worlds best microscope. BSM (H, top,  $\nu$ , SUSY..) Transformations: Searches at LHC, LHC as Higgs Precision Facility, QCD of Nuclear Dynamics  
The LHeC has a deep, unique QCD, H and BSM precision and discovery physics programme.

**Time**: Determined by the Large Hadron Collider (HL LHC needs till  $\sim 2040$  for  $3\text{ ab}^{-1}$ ) 1802.04317  
LHeC: Detector Installation in 2 years, earliest in LS4 (2030/31).  
HE LHC: re-use ERL. In between HL-HE, 10 years time of ERL Physics (laser,  $\gamma\gamma$ ..)  
Very long term: FCC-eh

<http://lhec.web.cern.ch>

**Challenges**: Demonstration of ERL Technology (high electron current, multi-turn)  
Design 3-beam IR for concurrent ep+pp operation, New Detector with Taggers - in 10 years.

**The LHeC is a great opportunity to sustain deep inelastic physics within future HEP.**  
The cost of an ep Higgs event is  $O(1/10)$  of that at any of the 4  $e^+e^-$  machines under consideration  
It can be done: the Linac is shorter than 2 miles and the time we have longer than HERA had.

**CERN and world HEP**: Vital to make the High Luminosity LHC programme a success.


# LHeC Prospects

- The ep interaction does not disturb pp, i.e. the LHC may become a twin collider, ep and pp operate concurrently and no luminosity loss is planned for pp. This requires a premounted eh detector which may then be inserted in 2 years.
- At LS4 (~2030) the heavy ion LHC operation ends and one may propose a different use of IP2 which currently houses ALICE.
- The electron beam energy ( $> 50$  GeV) and luminosity ( $O(10^{34}) \text{ cm}^{-2} \text{ s}^{-1}$ ) goals are derived from Higgs, top and BSM physics, also DIS itself ( $F_L$ , low  $x \sim 1/s$ ).
- The cost of the  $O(1)$  TeV ep collider is a small fraction of any other big project currently under discussion. The LHC determines the time frame. This may extend considerably if CERN moves to HE LHC in the forties.
- The ERL technology is being developed worldwide (Darmstadt, Cornell, Berlin, Novosibirsk, Jefferson Lab). PERLE would be a multi-turn 802 MHz ERL technology development and test facility which would timely accompany the LHeC progress.
- We celebrate this year the 50<sup>th</sup> anniversary of the discovery of quarks. This was not planned and achieved by a step in energy with a linac SHORTER than LHeC's
- There is a very long term future for eh as part of hh in the FCC vision

# Further use of ERL in between HL and HE LHC


## Reconfiguring LHeC → SAPPHiRE

### LHeC-ERL



### SAPPHiRE\*


### $\gamma\gamma$ Higgs factory




\*Small Accelerator for Photon-Photon Higgs production using Recirculating Electrons  
 S. A. Bogacz, J. Ellis, L. Lusito, D. Schulte, T. Takahashi, M. Velasco, M. Zanetti, F. Zimmermann,  
 'SAPPHiRE: a Small Gamma-Gamma Higgs Factory,' arXiv:1208.2827

F.Zimmermann at LHeC WS 9/17

## LHeC: perfect FCC-ee injector!



## LHeC-FEL



up to 60 GeV,  
 $\sim 25$  mA,  
 1 MeV photons?

3-15x higher beam energy  
 (10-200x higher  $\gamma$  energies),  
 300-600x higher current

XFEL: 20GeV e, 0.03mA, 24keV photons. LCLSII: 4 GeV e, 0.06mA, 5 keV photons

<b>4</b>	<b>An Electron Energy Recovery Linac at CERN</b>	<b>8</b>
4.1	The ERL Configuration of the LHeC . . . . .	9
4.1.1	Basic Structure and Design Choices . . . . .	9
4.1.2	Component Summary . . . . .	9
4.1.3	Performance Estimates and Operation Scenarios . . . . .	9
4.1.4	Electron-Ion Scattering . . . . .	9
4.2	ERL Beam Dynamics and Performance Limitations . . . . .	9
4.2.1	Front-end Simulation . . . . .	9
4.2.2	BBI, Filling Patterns and Bunch Structure . . . . .	9
4.3	LINAC and SRF . . . . .	9
4.3.1	Choice of Frequency and Cavity Prototype . . . . .	9
4.3.2	Sources . . . . .	9
4.3.3	Injector . . . . .	9
4.3.4	LINAC and Cavity-CryoModule . . . . .	9
4.3.5	Infrastructure: Power and Cryogenics . . . . .	9
4.4	Magnets and Dumps . . . . .	9
4.4.1	Arc Magnets . . . . .	9
4.4.2	Dumps . . . . .	9
4.5	Interaction Region . . . . .	9
4.5.1	Layout . . . . .	9
4.5.2	Triplet Magnet Design . . . . .	9
4.5.3	Optics . . . . .	9
4.5.4	IR and L* Optimisation . . . . .	9
4.6	Civil Engineering . . . . .	9
4.7	Stand-Alone Operation and Physics with the ERL, post HL LHC . . . . .	9
4.7.1	Photo-Nuclear Physics with the ERL . . . . .	9
4.7.2	A High Energy and Intensity Laser Facility . . . . .	9
4.7.3	A Photon-Photon Higgs Collider at CERN . . . . .	9
<b>5</b>	<b>Executive Summary and Further Project Development</b>	<b>10</b>
<b>6</b>	<b>Appendix</b>	<b>11</b>
6.1	The Technology Development Facility PERLE . . . . .	11
6.2	Cost-Energy-Physics Optimisation of the LHeC . . . . .	11

**DRAFT**



1	Modern Particle Physics and the Large Hadron Collider	4
2	Higgs, Discovery and Precision Physics	5
2.1	Electron-Hadron Scattering at the HL and HE LHC	6
2.1.1	Kinematics and Reconstruction of Final States	6
2.1.2	Opportunities through Energy Frontier ep and eA Interactions	6
2.1.3	Luminosity and Operation of an eh Collider at the LHC	6
2.2	Higgs Physics	6
2.2.1	High Precision Higgs Coupling Measurements	6
2.2.2	Htt Coupling Measurement	6
2.2.3	Combined ep and pp Analysis - the Potential for Precision Higgs Physics at the LHC	6
2.2.4	Discovery of the H-HH Selfcoupling	6
2.2.5	Exotic Higgs Decays and Dark Matter	6
2.2.6	Extension of the SM Higgs Sector	6
2.3	Challenging the Standard Model through High Precision and Energy	6
2.3.1	Resolving the Parton Substructure of the Proton	6
2.3.2	Electroweak, W Boson and Top Quark Physics	6
2.3.3	Non-Conventional Proton Structure Resolution	6
2.4	Searches for Physics Beyond the Standard Model	6
2.4.1	Heavy Neutrinos	6
2.4.2	Flavour Changing Neutral Currents	6
2.4.3	Substructure	6
2.4.4	Leptoquarks	6
2.5	Discovery through Precision QCD	6
2.5.1	High Mass and Bjorken x Searches at the LHC	6
2.5.2	Strong Coupling and Grand Unification	6
2.5.3	New QCD Dynamics at small x	6
2.5.4	A New Era of Nuclear Particle Physics with eA	6
3	Experimentation at the LHeC	7
3.1	Detector Design Considerations for HL and HE LHC	7
3.2	Detector Overview	7
3.2.1	Magnets	7
3.2.2	Interaction Region, Beam Pipe and Radiation	7
3.2.3	Inner Tracking	7
3.2.4	Calorimetry	7
3.2.5	Muon Detector	7
3.2.6	Central Detector Performance	7
3.3	Forward and Backward Detectors	7
3.3.1	Proton and Neutron Forward Taggers	7
3.3.2	Electron and Photon Backward Detectors	7
3.4	Detector Installation and Infrastructure	7

**DRAFT**

# Observations post CDR/EUSPP - 2012+ affecting ep at CERN:

**LHC lifetime** now extended to ~2040 to collect 3 [4]  $\text{ab}^{-1}$ . LS3 2024-2026+..

**Discovery of the Higgs:**  $L(\text{ep}): 10^{33} \rightarrow 10^{34} \text{ cm}^{-2} \text{ s}^{-1}$  [HERA in days]

**LHC brightness**  $N_p/\epsilon$  about 3 times higher than “ultimately” expected

**No further discovery at the LHC**, so far

**Detector technology** developments (LHC Det. Upgrades)

Strong **accelerator technology** developments, notably SCRF

ERL. LHeC: 720  $\rightarrow$  802 MHz. enhanced  $Q_0 > 10^{10}$

**EU strategy 13:** exploit LHC, study Higgs, develop SCRF+,

CERN: new accelerators “with emphasis on pp and ee”

Fine with the LHeC cost being a small fraction of ILC,CLIC,FCC

$\rightarrow$  CERN in 14 set up a new LHeC organisation with a new mandate and IAC (H.Schopper et al) to prepare for the next EU strategy 2019+

**Two main tasks (IAC): Update of CDR for HL-LHeC/ FCCeh + Testfacility**

# Framework of the Development

Following the CDR in 2012: Mandate issued by CERN:2014 (RH), confirmed in 2016 (FG)

## **Mandate to the International Advisory Committee**

Advice to the LHeC Coordination Group and the CERN directorate by following the development of options of an ep/eA collider at the LHC and at FCC, especially with:

Provision of scientific and technical direction for the physics potential of the ep/eA collider, both at LHC and at FCC, as a function of the machine parameters and of a realistic detector design, as well as for the design and possible approval of an ERL test facility at CERN.

Assistance in building the international case for the accelerator and detector developments as well as guidance to the resource, infrastructure and science policy aspects of the ep/eA collider.

Chair: Herwig Schopper, em. DG of CERN. IAC+CERN have invited four of its members to follow the study with special attention (Stefano Forte, Andrew Hutton, Leandro Nisati and Lenny Rifkin). Collaboration also with the FCC Review Committee chaired by Guenther Dissertori.

LHeC has been a development for and initiated by CERN, ECFA and NuPECC, so far, it's formal status is that of a community study, not a proposal, which holds for the FCC also, of which 'eh' is a part.

# Organisation\*)

## International Advisory Committee

Mandate by CERN to define

“..Direction for ep/A both at LHC+FCC”

Sergio Bertolucci (CERN/Bologna)

Nichola Bianchi (Frascati)

Frederick Bordry (CERN)

Stan Brodsky (SLAC)

Hesheng Chen (IHEP Beijing)

Eckhard Elsen (CERN)

Stefano Forte (Milano)

Andrew Hutton (Jefferson Lab)

Young-Kee Kim (Chicago)

Victor A Matveev (JINR Dubna)

Shin-Ichi Kurokawa (Tsukuba)

Leandro Nisati (Rome)

Leonid Rivkin (Lausanne)

**Herwig Schopper (CERN) – Chair**

Jurgen Schukraft (CERN)

Achille Stocchi (LAL Orsay)

John Womersley (ESS)

**We miss Guido Altarelli.**

## Coordination Group

Accelerator+Detector+Physics

Gianluigi Arduini

Nestor Armesto

Oliver Brüning – Co-Chair

Andrea Gaddi

Erk Jensen

Walid Kaabi

Max Klein – Co-Chair

Peter Kostka

Bruce Mellado

Paul Newman

Daniel Schulte

Frank Zimmermann

**5(12) are members of the  
FCC coordination team**

OB+MK: co-coordinate FCCeh

## Working Groups

### PDFs, QCD

Fred Olness,

Claire Gwenlan

### Higgs

Uta Klein,

Masahiro Kuze

### BSM

Georges Azuelos,

Monica D’Onofrio

Oliver Fischer

### Top

Olaf Behnke,

Christian

Schwanenberger

### eA Physics

Nestor Armesto

### Small x

Paul Newman,

Anna Stasto

### Detector





Alessandro Polini

Peter Kostka

\*) April 2018



# 60 GeV Energy Recovery Linac



CDR: Default configuration, 60 GeV, 3 passes, 720 MHz, synchronous ep+pp,  $L_{ep}=10^{33}$



