

Vector calculus

- In this lecture we will:
 - ◆ Sketch out how we can derive a potential from a field using line integrals.
 - ◆ Do an example to check it works!
 - ◆ Look at a physical example: deriving the electric potential from the electric field.
 - ◆ Mention a caveat: there are some fields that cannot be derived from potentials.
- Some comprehension questions for this lecture.
 - ◆ What is the potential associated with the field:

$$\vec{E}(x, y, z) = \begin{pmatrix} yz + 2xy \\ x^2 + xz + z^2 \\ xy + 2yz \end{pmatrix} ?$$


1

Deriving a potential from a field

- We have seen that we can get a field from a potential: $\vec{F}(x, y, z) = \nabla\phi(x, y, z)$.
- Suppose we have a field $\vec{F}(x, y)$, can we derive from this the associated potential $\phi(x, y)$?
- Illustrate idea in 2D (more formal proof in text books!).
- Consider stepping from A to B in the scalar field $\phi(x, y)$.
- Change in ϕ is $\delta\phi$, given by slope in direction of movement and step length.
- For step δx in x direction:

$$\delta\phi_x \approx \frac{\partial\phi(x, y)}{\partial x} \delta x = F_x \delta x.$$
- For subsequent step δy in y direction

$$\delta\phi_y \approx \frac{\partial\phi(x + \delta x, y)}{\partial y} \delta y \approx \frac{\partial\phi(x, y)}{\partial y} \delta y \approx F_y \delta y.$$
- If step in x then y, $\delta\phi \approx \delta\phi_x + \delta\phi_y$.



2

1

2

Deriving a potential from a field

- Rewriting this:


$$\delta\phi \approx F_x \delta x + F_y \delta y.$$
- Now take n steps from initial position i to final position f:
 - The total change in ϕ is then

$$\sum_n \delta\phi_n \approx \sum_n F_x(x_n, y_n) \delta x_n + F_y(x_n, y_n) \delta y_n$$
 - Taking the limit of infinitely many infinitely small steps:

$$\int d\phi = \int_C F_x(x, y) dx + F_y(x, y) dy$$

$$\phi = \phi(x_i, y_i) + \int_C (F_x, F_y) \cdot (dx, dy)$$

$$= \phi(x_i, y_i) + \int_C \vec{F} \cdot d\vec{r}$$
 - The subscript C tells us to move along curve from i to f.
 - If start at (0, 0) and move to (x, y) we have “climbed” $\phi(x, y) - \phi(0, 0)$.



3

Deriving a potential from a field

- Example:
 - Field $\vec{E}(x, y) = (2x + y, x)$.
 - Find the associated potential,

$$\phi = \phi_0 + \int_C \vec{E} \cdot d\vec{r}$$
 - Integrate along $\vec{r}(t) = (x(t), y(t)) = (xt, yt), t = 0 \dots 1$.

$$\phi = \phi_0 + \int_0^1 \vec{E}(x(t), y(t)) \cdot \frac{d\vec{r}(t)}{dt} dt$$

$$= \phi_0 + \int_0^1 E_x(x(t), y(t)) \frac{dx(t)}{dt} dt + \int_0^1 E_y(x(t), y(t)) \frac{dy(t)}{dt} dt$$
 - Then:

$$\phi = \phi_0 + \int_0^1 [2(xt) + (yt)] \times x dt + \int_0^1 (xt) \times y dt$$

$$= \phi_0 + \frac{t^2}{2} \times (2x^2 + xy) + \frac{t^2}{2} \times xy \Big|_0^1$$

$$= \phi_0 + x^2 + \frac{xy}{2} + \frac{xy}{2}$$

$$= \phi_0 + x^2 + xy.$$
 - Check:

$$\nabla\phi = \begin{pmatrix} \frac{\partial}{\partial x} x^2 + xy + \phi_0 \\ \frac{\partial}{\partial y} x^2 + xy + \phi_0 \end{pmatrix} = \begin{pmatrix} 2x + y \\ x \end{pmatrix} = \vec{E}$$

4

3


4

Electric potential from electric field

- Electric field due to point charge given by:

$$\vec{E} = \frac{q}{4\pi\epsilon_0} \begin{pmatrix} \frac{x}{(x^2 + y^2 + z^2)^{\frac{3}{2}}} \\ \frac{y}{(x^2 + y^2 + z^2)^{\frac{3}{2}}} \\ \frac{z}{(x^2 + y^2 + z^2)^{\frac{3}{2}}} \end{pmatrix}$$

- Write $\frac{q}{4\pi\epsilon_0} = K$.



- Using line integral method:

$$\begin{aligned} \phi &= \phi_0 + \int_c \vec{E} \cdot d\vec{r} \\ &= \phi_0 + \int_0^1 E_x \frac{dx}{dt} dt + \int_0^1 E_y \frac{dy}{dt} dt + \int_0^1 E_z \frac{dz}{dt} dt \end{aligned}$$

$$\text{with } \vec{r}(t) = \begin{pmatrix} xt \\ yt \\ zt \end{pmatrix} \text{ and taking the path}$$

to be from $t=0$ to $t=1$ as before.

5

Electric potential from electric field

- Look at E_x integral:

$$\begin{aligned} \int_0^1 E_x \frac{dx}{dt} dt &= K \int_0^1 \frac{xt}{((xt)^2 + (yt)^2 + (zt)^2)^{\frac{3}{2}}} x dt \\ &= K \frac{x^2}{(x^2 + y^2 + z^2)^{\frac{3}{2}}} \int_0^1 \frac{t}{(t^2)^{\frac{3}{2}}} dt \\ &= K \frac{x^2}{(x^2 + y^2 + z^2)^{\frac{3}{2}}} \int_0^1 \frac{1}{t^2} dt \\ &= K \frac{x^2}{(x^2 + y^2 + z^2)^{\frac{3}{2}}} \left. \frac{-1}{t} \right|_0^1 \end{aligned}$$

- There is a problem, can't evaluate one limit of integral: \vec{E} infinite at origin!

- One solution is to change the path.

- Move from point at infinity to position (x, y, z) then have:

$$\begin{aligned} \int_{\infty}^1 E_x \frac{dx}{dt} dt &= K \frac{x^2}{(x^2 + y^2 + z^2)^{\frac{3}{2}}} \left. \frac{-1}{t} \right|_{\infty}^1 \\ &= -K \frac{x^2}{(x^2 + y^2 + z^2)^{\frac{3}{2}}} \end{aligned}$$

- Repeat for E_y and E_z and add results:

$$\phi = \phi_{\infty} - K \frac{(x^2 + y^2 + z^2)}{(x^2 + y^2 + z^2)^{\frac{3}{2}}} = -\frac{q}{4\pi\epsilon_0} \frac{1}{r}$$

- Note minus sign, not present when "physics convention" used, we have decided $\vec{E} = -\nabla\phi$ not $\vec{E} = \nabla\phi$.

6

Caveat: fields that are not derivable from potentials

- Recall from lecture 6: $\nabla \times (\nabla\phi) = 0$.

- Hence, a vector field derived from a potential, e.g. $\vec{E} = \nabla\phi$, must always satisfy $\nabla \times \vec{E} = 0$.

- Conversely, a field for which the curl is not zero cannot be derived from a potential.

E.g. $\vec{F} = \begin{pmatrix} yz \\ xz - 3yz \\ xy + 2z \end{pmatrix}$.

- Using our prescription...

$$\begin{aligned} \phi &= \phi_0 + \int_0^1 yt zt x dt + \int_0^1 (xt zt - 3yt zt) y dt \\ &\quad + \int_0^1 (xt yt + 2zt) z dt. \end{aligned}$$

- So: $\phi = \phi_0 + xyz \int_0^1 t^2 dt + (xyz - 3y^2z) \int_0^1 t^2 dt$

$$+ xyz \int_0^1 t^2 dt + 2z^2 \int_0^1 t dt$$

$$= \phi_0 + xyz \frac{t^3}{3} + (xyz - 3y^2z) \frac{t^3}{3} \Big|_0^1$$

$$+ xyz \frac{t^3}{3} + 2z^2 \frac{t^2}{2} \Big|_0^1$$

$$\Rightarrow \phi = \phi_0 + xyz - y^2z + z^2$$

- Now calculate field:

$$\nabla(xyz - y^2z + z^2) = \begin{pmatrix} yz \\ xz - 2yz \\ xy - y^2 + 2z \end{pmatrix} \neq \vec{F}$$

7