

# HERA Collider Physics

Durham, 6.12.2001

*Peter Schleper*

*DESY*

**HERA results in 2006**

a personal selection

incomplete


speculative



# $e^\pm p$ collisions at HERA I and II


## HERA I: 1993-2000

- $E_{p_0} = 820 \dots 920 \text{ GeV}$
- $E_{e_0} = 27.5 \text{ GeV}$
- $\sqrt{s} = 320 \text{ GeV}$
- $L \gtrsim 100 \text{ pb}^{-1}$




## HERA II: 2001-2006 (?)

- Increase to  $L=1000 \text{ pb}^{-1}$
- $e^+$  and  $e^-$  equally shared
- $e^\pm$  polarisation  $\approx 55\%$
- Runs with lower/higher  $E_{p_0}$



# HERA II



## Collider experiments: H1, ZEUS

- Stronger focussing close to experiments
- Spin rotators
- New beam lines

## HERMES, HERA-B


- not affected

# HERA II

---

## The HERA upgrade

- Strong focussing at interaction point
- Magnets inside main detector



## The Challenges

- Synchrotron radiation
- Back scattering from magnets

## The Detector upgrades

- Trigger !
- New forward tracking
- Extended silicon tracking
- New Forward proton spectrometer
- Smaller acceptance at low angles:  $Q^2 > 8 \text{ GeV}^2$

# Virtues of $ep$ scattering at HERA

## large $Q^2$ lever arm: precision QCD

- $\alpha_s$ , quark and gluon densities

## small - $x$ : high parton densities

- novel quantum system

## small $Q^2$ : large distance QCD


- non-perturbative, confinement

## large CMS energy:

- (beyond) the EW standard model

(largest  $E_{CMS}$  with lepton in initial state)

→ reason why H1 / ZEUS are leading QCD experiments



# Partons in the Proton

---


$ep \rightarrow eX$  at small  $Q^2$ :

$$\frac{d^2\sigma}{dx dQ^2} \approx \frac{2\pi\alpha^2}{xQ^4} F_2$$

momentum distribution of quarks:

$$F_2(x, Q^2) \approx \sum e_q^2 q(x, Q^2)$$

$$dF_2/d\ln Q^2|_x \approx \alpha_s \times g(x, Q^2)$$





small  $x \rightarrow$  high quark density

$\rightarrow$  strong rise of quark density  $\rightarrow$  high gluon density

---

# Quark/Gluon Distribution in the Proton




# Higgs Produktion at LHC

---

$M_H > 108 \text{ GeV}$

LHC parton kinematics



depends on gluon density at  $x \lesssim 10^{-3}$


only process: DIS at HERA

but:  $x \lesssim 10^{-4} \rightarrow Q^2 \lesssim 1 \text{ GeV}$

# Gluon induced processes


---

Verification of gluon density: intermediate  $x$



jet production

H1 Collaboration



charm production


H1 Collaboration

# Longitudinal Structure Function $F_L$

Verification of gluon density: small  $x$


$$F_L \sim \sigma_r - F_{2,\text{extrapol.}}$$

- depends on assumption for extrapolation towards high  $y$
- limited in precision
- includes small  $x$



$F_L$  from reduced beam energy


- e.g.  $E_P = 300, 350, 465$  GeV




**X**

# Bottom production: Unexplained


HERA now



CDF




HERA II




much enhanced B-tagging

# Tevatron jets at high $P_T$ : gluons at high $x$

CDF: cone algorithm



D0:  $K_T$  algorithm




open questions...


# Photon and Proton at High x

---

jets in  $\gamma p$  at high  $E_T$




access to proton at high x !



- dominated by direct + pointlike part of  $\gamma$
- significant contribution from gluon in proton  
→ Tevatron high  $E_T$  jets

## HERA summary on $\alpha_s$

---



- NNLO calculations for F2 (and 2-jet production ?)

→ reduction of scale errors by factor 2-3 !!


the CHALLENGE: get best  $\alpha_s$  ( $\pm 1\%$ )

needs reduction also of experimental errors !!


---

# Low- $x$ behaviour of $F_2$


$$\lambda =: - \left( \frac{d \ln F_2}{d \ln x} \right)_{Q^2}$$




- no sign of deviation from  $x^{-\lambda(Q^2)}$  at small  $x$
- no saturation visible in  $F_2$



# Low- $x$ Parton Dynamics




- small- $x$ , forward jet ( $\pi^0$ ),  
 $Q^2 \approx E_{T,jet}^2$
- no  $k_T$  order, test case for  
DGLAP–BFKL–CCFM– $\gamma^*$ <sub>resolved</sub>
- unintegrated parton distributions



## No direct dynamical evidence up to now

- CCFM (CASCADE): some success for  $F_2^{charm}$
- NNLO for jet production needed

# Hard Diffraction



colour singlet exchange of more than one parton (2 gluons or more)  
→ Correlations

## Hard scattering factorization

$$\sigma \sim \sum_i f_{i/p}(x_F, t, x, Q^2) \times \hat{\sigma}_i(x, Q^2)$$

with DGLAP evol.  $f_{i/p}(x_F, t, x, Q^2)$  at fixed  $x_F, t$ .

- Proof for diffractive DIS
- Experimental test ?
- Partonic picture of diffraction



## Colour Dipol Picture of DIS

- hope to unify description of DIS and Diffraction

## GPD's: Generalized Parton Distributions

- $f_{i/p}(x_1, x_2, Q^2)$

# Diffractive Structure Functions



limited by systematics

limited by statistics

# Very Forward Proton Spectrometer (VFPS)


New detector at  $z = 200m$

- full acceptance at  
 $0.005 < x_{IP} < 0.02$ ,  
 $|t| < 0.25 \text{ GeV}^2$
- trigger for all diffractive processes
- $|t|, \phi$  measurement
- no contribution from p-dissociation

→ High statistics test of QCD  
interpretation of diffraction:  $F_2^{D(4)}$ ,  
jets, charm, DVCS

H1 VFPS acceptance


Beam pipe aperture acceptance



# Colour Dipol Picture of DIS

Assume factorisation:

$$\sigma = \Psi(\gamma^* \rightarrow qq(g)) \times \sigma_{\text{dipol}}$$




## Success of Colour Dipol Models

- fit  $\sigma_{\text{dipol}}$  to  $F_2$
- predict  $F_2^D$ , diffractive  $J\Psi$ , jets
- Connection to BFKL

$Q^2 \rightarrow 0$ : saturation of  $\sigma_{\text{dipol}}$



(Golec-Biernat, Wüsthoff)

- not yet successful at  $Q^2 < 1 \text{ GeV}^2$



# Deeply Virtual Compton Scattering

---



$\gamma$ - × gluon-pair → QCD describes data !

access to parton correlations: “skewed pdf’s”

→ new forward proton tagger,  $e_{L,R}^\pm$

---

# DIS cross section

---

## Neutral Current

$$\frac{d^2\sigma_{NC}^{e\pm p}}{dx dQ^2} = \frac{2\pi\alpha^2}{xQ^4} \left[ Y_+ \tilde{F}_2(x, Q^2) \mp Y_- x \tilde{F}_3(x, Q^2) - y^2 \tilde{F}_L(x, Q^2) \right]$$

$$Y_\pm \equiv 1 \pm (1-y)^2$$

$$\tilde{F}_2 \equiv F_2 - v_e \frac{\kappa_w Q^2}{(Q^2 + M_Z^2)} F_2^{\gamma Z} + (v_e^2 + a_e^2) \left( \frac{\kappa_w Q^2}{Q^2 + M_Z^2} \right)^2 F_2^Z = x \sum A_i (q_i + \bar{q}_i)$$

$$x \tilde{F}_3 \equiv - a_e \frac{\kappa_w Q^2}{(Q^2 + M_Z^2)} x F_3^{\gamma Z} + (2v_e a_e) \left( \frac{\kappa_w Q^2}{Q^2 + M_Z^2} \right)^2 x F_3^Z = x \sum B_i (q_i - \bar{q}_i)$$

$v_e, a_e$  - vector and axial couplings,  $\kappa_w^{-1} = 4 \frac{M_W^2}{M_Z^2} (1 - \frac{M_W^2}{M_Z^2})$

## Charged Current

$$\frac{d^2\sigma_{CC}^\pm}{dx dQ^2} = \frac{G_F^2 M_W^4}{2\pi x} \frac{1}{(Q^2 + M_W^2)^2} \phi_{CC}^\pm(x, Q^2)$$

$$\phi_{CC}^+ = x [(\bar{u} + \bar{c}) + (1-y)^2(d+s)] \quad (in LO)$$

$$\phi_{CC}^- = x [(u+c) + (1-y)^2(\bar{d}+\bar{s})]$$


## Reduced NC and CC cross sections

$$\tilde{\sigma}_{NC}(x, Q^2) \equiv \frac{1}{Y_+} \frac{Q^4}{2\pi\alpha^2} x \frac{d^2\sigma_{NC}}{dx dQ^2} = F_2 (1 + \Delta_{F_2} + \Delta_{F_3} + \Delta_{F_L})$$

$$\tilde{\sigma}_{CC}(x, Q^2) \equiv \frac{2\pi x}{G_F^2} \left( \frac{M_W^2 + Q^2}{M_W^2} \right)^2 \frac{d^2\sigma_{CC}}{dx dQ^2}$$



---

# High $Q^2$ and Weak Interaction




$$\gamma/Z \approx W$$

Unification



needs more data at large  
 $x, Q^2$  !

## Cross sections at high $Q^2$




$\gamma/Z$ : mainly  $u$ - quarks

## Cross sections at high $Q^2$

---

b




$\gamma/Z$ : mainly  $u$ - quarks


---

# Cross sections at high $Q^2$


$xF_3$ : Valence quarks



$W$ -exchange: mainly  $d$ - quarks:  $\pm$




# Polarized NC Cross Sections



## $\gamma Z$ interference

- $\gamma^2 \approx \gamma Z \approx Z^2$  for  $\frac{\kappa Q^2}{Q^2 + M_Z^2} = 1$   
 $\rightarrow Q^2 \approx 2M_Z^2 \approx 16000 \text{ GeV}^2$
- positive / negative interference for  $e^- p / e^+ p$
- 10% effect at high  $Q^2$


# Weak couplings of light quarks



- needs measurement of  $e_L^+, e_L^-, e_R^+, e_R^-$
- only light quarks from proton
- similar precision as from LEP

# Right handed weak currents: $W_R$

---




## Polarized charged currents


- $\sigma_{CC}(e_P^\pm) = (1 \pm P) \sigma_{CC}(e_{P=0}^\pm)$
- needs high polarisation
- needs high precision of polarisation
- $M_{W_R} \lesssim 600$  GeV
- similar to direct search at Tevatron

# Beyond the Standard Model: Supersymmetrie

single production of squarks in  $R$ -parity violating models



## Minimal Supergravity + $R_p$ Violation



- nothing found yet up to  $M_{\tilde{q}} = 245$  GeV
- will reach sensitivity of  $M_{\tilde{q}} \approx 300$  GeV

# Summary

---

What did / will we learn from HERA ?

- low-x: new regime of QCD
  - the main theoretical activity so far
  - QCD picture of Diffraction
  - colour dipole — BFKL ??
- precision QCD:
  - world best value of  $\alpha_s$
  - Final (?) word on proton structure at HIGH and LOW  $x$
- high  $Q^2$ : only just started
  - luminosity upgrade
  - precision test of SM at  $10^{-18}$  m
- searches:
  - high  $P_T$  lepton events ?
  - Supersymmetry with  $R_P$  violation
  - contact interactions
  - + ??