
#### Vector calculus

- In this lecture we will:
  - Define the divergence of a vector field.
  - Look at some examples to try and gain some insight into what the divergence represents.
  - Discuss the divergence of the electric and magnetic fields.

- Some comprehension questions for this lecture.
  - ❖ Indicate where the divergence will be positive below.



• Calculate the divergence of the field  $\vec{F}(x, y, z) = (x^2 \ 2xy \ xyz)$ .

The divergence of a vector field is defined by the expression:

$$\nabla \cdot \vec{F}(x, y, z) = \frac{\partial}{\partial x} F_x + \frac{\partial}{\partial y} F_y + \frac{\partial}{\partial z} F_z$$


- Note, the divergence of a vector field is a scalar field.
- Can think of the divergence as dot product of vector with the operator

$$\nabla = \left( \frac{\partial}{\partial x} \quad \frac{\partial}{\partial y} \quad \frac{\partial}{\partial z} \right).$$

Look at an example (in 2D so can visualise more easily):

$$\vec{F}(x,y) = \left(\frac{x}{\sqrt{x^2 + y^2}} \quad \frac{y}{\sqrt{x^2 + y^2}}\right)$$

Plot vector field  $\bar{F}(x, y)$ :



Calculate divergence, x component... ...and y component


$$\begin{split} \frac{\partial}{\partial x} F_x &= \frac{\partial}{\partial x} \frac{x}{\sqrt{x^2 + y^2}} \\ &= \frac{\sqrt{x^2 + y^2} - x \times \frac{1}{2} (x^2 + y^2)^{-\frac{1}{2}} 2x}{x^2 + y^2} \\ &= \frac{1}{\sqrt{x^2 + y^2}} - \frac{x^2}{\left(x^2 + y^2\right)\sqrt{x^2 + y^2}} \\ &= \frac{x^2 + y^2 - x^2}{\left(x^2 + y^2\right)\sqrt{x^2 + y^2}} \\ &= \frac{y^2}{\left(x^2 + y^2\right)^{\frac{3}{2}}} \end{split}$$

$$\frac{\partial}{\partial y} F_y = \frac{\partial}{\partial y} \frac{y}{\sqrt{x^2 + y^2}}$$
$$= \frac{x^2}{\left(x^2 + y^2\right)^{\frac{3}{2}}}$$


Putting these together:

$$\nabla \cdot \vec{F} = \frac{y^2}{\left(x^2 + y^2\right)^{\frac{3}{2}}} + \frac{x^2}{\left(x^2 + y^2\right)^{\frac{3}{2}}}$$
$$= \frac{1}{\sqrt{x^2 + y^2}} = \frac{1}{r}$$


#### Plot $\vec{F}$ and $\nabla \cdot \vec{F}$ :



#### Plot $\vec{G}(x, y) = -\vec{F}(x, y)$ and $\nabla \cdot \vec{G}$ :



Divergence of field reveals "sources" (left) and "sinks" (right) of field.




### Divergence of a field

- A field is defined by the equation:  $\vec{F}(x, y, z) = (3\sin x + 2\cos x - z^2).$
- Determine the divergence of the field.
- What is the value of  $\nabla \cdot \vec{F}(\frac{\pi}{2}, 0, -1)$ ?

- Look for sources and sinks in another  $\blacksquare$  Plot  $\vec{F}$  and  $\nabla \cdot \vec{F}$ : vector field:
- $\vec{F}(x,y) = \begin{pmatrix} \sin 2x \\ \cos 2y \end{pmatrix}$
- Divergence of this field is:


$$\nabla \cdot \vec{F}(x, y) = \frac{\partial}{\partial x} \sin 2x + \frac{\partial}{\partial y} \cos 2y$$
$$= 2\cos 2x - 2\sin 2y$$



- Same as previous slide using contours to represent divergence.
- $\vec{F}(x,y) = \begin{pmatrix} \sin 2x \\ \cos 2y \end{pmatrix}$
- Divergence is:

$$\nabla \cdot \vec{F}(x, y) = \frac{\partial}{\partial x} \sin 2x + \frac{\partial}{\partial y} \cos 2y$$
$$= 2\cos 2x - 2\sin 2y$$

#### Plot $\vec{F}$ and $\nabla \cdot \vec{F}$ :




#### Divergence of electric field

One of Maxwell's equations (Gauss' Law) involves divergence of E field:

$$\nabla \cdot \vec{E} = \frac{\rho}{\epsilon_0}$$


- The quantity ρ is the density of electric charge.
- This equation is telling as that electric charge is the source of the electric field.



### Divergence of electric field

- Another of Maxwell's equations (Gauss' Law for magnetism) involves the divergence of the magnetic field:  $\nabla \cdot \vec{B} = 0$
- What does this equation tell us about the sources and sinks of the magnetic field?
- And about magnetic monopoles?

Example of how magnetic field can be generated:



- How is magnetism caused in materials, e.g. "magnets"?
- In the earth?
- Reversal of Earth's magnetic field.